

Acute HIV infection in a 39-year-old man

Shinya Matsumoto MD PhD, Yuki Murata MD, Yoshitaka Tomoda MD PhD

■ Cite as: *CMAJ* 2022 November 21;194:E1541. doi: 10.1503/cmaj.220656

A 39-year-old man presented to the emergency department with a 7-day history of fever and rash. He reported same-sex relationships involving condomless anal sex, and did not use pre- or postexposure HIV prophylaxis. Physical examination found a temperature of 38°C, cervical lymphadenopathy, mucosal ulceration on his tongue and a maculopapular rash on his face, neck and anterior chest (Figure 1). Suspecting primary HIV infection, we performed a rapid plasma reagin test, which returned a negative result. A chemiluminescent immunoassay was positive for HIV antibodies, but the reactive immunoassay using Western blotting showed no reactivity. The patient's HIV-1 blood viral load was 4.79 million copies/mL with a blood CD4 T-cell count of $0.1 \times 10^9/L$. We diagnosed acute HIV infection and started antiretroviral treatment with bictegravir–emtricitabine–tenofovir alafenamide.

Features of acute HIV infection occur 2–4 weeks after exposure; seroconversion usually occurs 3–8 weeks after initial infection. Common presentation includes fever, arthralgia and lymphadenopathy, which may mimic acute mononucleosis.¹ Mucocutaneous ulceration is highly suggestive of acute HIV infection.² Rash occurs in about 50% of patients, typically 3 days after fever onset, and persists for 5–8 days. It is characterized by small (5–10 mm), well-circumscribed, erythematous macules or maculopapules, mainly on the anterior chest.^{1,2} A history of high-risk sexual contact raises suspicion of acute HIV infection.

Transmission rates among patients with acute HIV infection are 9–15 times greater than among those with chronic infections.³ However, the diagnosis of acute HIV infection is often missed owing to the asymptomatic or self-limited nature of the virus, nonspecific signs (such as the rash seen in our patient) or clinician hesitancy to ask questions about sexual exposure. Immediate initiation of antiretroviral therapy in the early stage of HIV infection reduces viral load.⁴

References

1. Chu C, Selwyn PA. Diagnosis and initial management of acute HIV infection. *Am Fam Physician* 2010;81:1239-44.
2. Richey LE, Halperin J. Acute human immunodeficiency virus infection. *Am J Med Sci* 2013;345:136-42.




Figure 1: Photographs of a 39-year-old man with acute HIV infection, showing (A) small, well-circumscribed, red maculopapular rashes (arrows) on his anterior thorax. (B) Close-up image of the rash.

3. Hollingsworth TD, Pilcher CD, Hecht FM, et al. High transmissibility during early HIV infection among men who have sex with men—San Francisco, California. *J Infect Dis* 2015;211:1757-60.
4. The INSIGHT START Study Group; Lundgren JD, Babiker AG, Gordin F, et al. Initiation of antiretroviral therapy in early asymptomatic HIV infection. *N Engl J Med* 2015;373:795-807.

Competing interests: None declared.

This article has been peer reviewed.

The authors have obtained patient consent.

Affiliation: Department of General Medicine, Itabashi Chuo Medical Center, Tokyo, Japan

Content licence: This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY-NC-ND 4.0) licence, which permits use, distribution and reproduction in any medium, provided that the original publication is properly cited, the use is noncommercial (i.e., research or educational use), and no modifications or adaptations are made. See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Correspondence to: Yoshitaka Tomoda, yoshisoph@gmail.com

Clinical images are chosen because they are particularly intriguing, classic or dramatic. Submissions of clear, appropriately labelled high-resolution images must be accompanied by a figure caption. A brief explanation (300 words maximum) of the educational importance of the images with minimal references is required. The patient's written consent for publication must be obtained before submission.