physician fees cannot decline in real terms indefinitely.

More important, the authors' data do not penetrate the effects of financial restraint on quality of care, a fact that they themselves point out.¹ But this issue is the very crux of the perceived health care crisis. What happened to waiting lists for referrals, surgery and diagnostic tests? How were health outcomes affected? Is it appropriate to assume that age- and population-adjusted fee expenditures *should* remain the same (in real dollars) over this time period? Moreover, physician fees account for only 24% of health care spending in British Columbia;² if there is a funding crisis, physician fees are only a small part of a larger problem.

A sustainable system must both control costs and provide appropriate health care. Barer and colleagues establish that the government of British Columbia controlled costs, but they do not establish that it did so in a sustainable manner.

Mark Fruitman

Radiologist

St. Joseph's Health Centre
Toronto, Ont.

References

1. Barer ML, Evans RG, McGrail KM, Green B, Hertzman C, Sheps SB. Beneath the calm surface: the changing face of physician-service use in British Columbia, 1985/86 versus 1996/97. *CMAJ* 2004;170(5):803-7.
2. Resource summary. In: *Budget 2004*. Service plan 2004/05-2006/07. Victoria: Ministry of Health Services; [date unknown]. Available: www.bcbudget.gov.bc.ca/sp2004/hs/hs.pdf (accessed 2004 Apr 5). p. 12 (2003/04 restated estimates).

DOI:10.1053/cmaj.1040426

[Three of the authors respond:]

We strongly support Mark Fruitman's point that "[a] sustainable system must both control costs and provide appropriate health care." Our purpose¹ was to identify the key components in physician billings as a means of isolating both what was and what was not happening. The observation of major changes in the patterns of contact between patients and general practitioners is, we think, quite new: we found that GPs are increasingly shar-

ing their patients, yet the net impact on costs is minimal. It appears that neither patients nor physicians are abusing the system or, if they are, that abuse has not been increasing. This situation might raise concerns about continuity of care, but such concerns take us beyond these data.

In contrast, the care of elderly patients after they pass the GP "gatekeeper" and enter the specialty system is much more expensive. As Fruitman points out, our data cannot say whether these dramatic increases in cost are appropriate. But if one is truly concerned about both the appropriateness and the cost of care, these sectors and this patient group would seem obvious places for further scrutiny.

Fruitman suggests that our conclusion regarding sustainability had its basis in costs for physicians, whose real fees declined over the period in question. But even if fees had increased at the rate of inflation, the average annual increase in expenditures would have been 6.8% rather than 5.8%, hardly enough difference to support claims of "unsustainability." Furthermore, this calculation presumes that the change in (age-specific) utilization per capita would have been the same, irrespective of the change in real fees. It seems conceivable that faster growth in fees would instead have been associated with slower growth in utilization — the fact that the increase in use almost precisely offset the decline in fees may be more than coincidence. In any case, physician fees in British Columbia have been the highest, or among the highest, in the country for decades.

Morris L. Barer

Robert G. Evans

Kimberlyn McGrail

Centre for Health Services and Policy
Research
University of British Columbia
Vancouver, BC

Reference

1. Barer ML, Evans RG, McGrail KM, Green B, Hertzman C, Sheps SB. Beneath the calm surface: the changing face of physician-service use in British Columbia, 1985/86 versus 1996/97. *CMAJ* 2004;170(5):803-7.

DOI:10.1053/cmaj.1040685

Reducing adverse events

Further to the landmark work of Alan Forster and coauthors¹ at the Ottawa Hospital–Civic Campus in identifying levels of adverse events at discharge and revealing the extent of the gap in continuity of care, particularly drug management, I wish to point out measures that have been taken in this region to diminish the problem.

In 1995 a group of Ottawa-area health care professionals — hospital and community pharmacists, physicians and a nurse — gathered to promote solutions to the gaps in seamless care.² We recommended 2 major innovations: pocket drug profiles to be used by community pharmacists for patients receiving long-term medication, as well as formal discharge communications from the hospital to the family physician and the community pharmacist. The latter recommendation was instituted the next year at the Ottawa General Hospital (now the Ottawa Hospital–General Campus) in a form called "Prescription and Discharge Notes," which provided complete information on discontinued medications, medication incidents and recommendations for ongoing care. A pilot project was instituted on a surgical floor, and a chart review followed. The review indicated that on average half of each patient's medications were changed before discharge and that potentially 61% of the discharge forms reported drug-related problems, only half of which were resolved before discharge.³ A later study on a medical floor showed an even greater number of changes and potential drug-related problems.⁴

This form has now been updated and its use extended to the Civic Campus and to other institutions. A further study showed that use of the form on the medical ward of a Montreal hospital increased the accuracy of patient profiles maintained by community pharmacists.⁵ It would be interesting to determine if this form makes a difference in the negative outcomes that Forster and coauthors¹ so clearly identified.

Many other communities across Canada have tackled this problem, as revealed by posters and presentations at 2 national seamless care workshops.^{6,7} The report by Forster and coauthors clearly demonstrates the need for pharmacists, physicians and others to mend the gap in communications between institutional and community care.

William McLean

Pharmaceutical Outcomes Research Unit
Ottawa Hospital — General Campus
Ottawa, Ont.

References

1. Forster AJ, Clark HD, Menard A, Dupuis N, Chernish R, Chandok N, et al. Adverse events among medical patients after discharge from hospital. *CMAJ* 2004;170(3):345-9.
2. Sullivan M, McLean W. Seamless care: a review of the issues. *Can Pharm J* 2000;133(7):26-9.
3. McLean W. The influence of a prescription and discharge notes form on drug-related problems [abstract]. American College of Clinical Pharmacy annual meeting; 1997 Nov 10; Phoenix.
4. Rogers K, Tierney M, Singh A, McLean W. Assessment of a seamless care prescription/discharge notes form. *Can J Hosp Pharm* 2003;56:14-23.
5. Paquette-Lamontagne N, McLean W, Besse L, Cusson J. Evaluation of a new integrated discharge prescription form. *Ann Pharmacother* 2001;35:953-8.
6. Proceedings of the seamless care workshop 1998; 1998 Oct 30 – Nov 1; Toronto. Ottawa: Canadian Society of Hospital Pharmacists, Canadian Pharmacists Association; 1999.
7. Proceedings of the seamless care workshop 2000; 2000 Sep 30; Val-Morin (QC). Ottawa: Canadian Society of Hospital Pharmacists, Canadian Pharmacists Association; 2001.

Competing interests: None declared.

DOI:10.1053/cmaj.1040362

[Two of the authors respond:]

The form to which William MacLean refers in his letter was being used during our study,¹ and we commonly saw problems with how the forms were completed. For example, it was often difficult to distinguish old from new medications, poor handwriting often made prescriptions illegible, and only rarely was a contact number for the prescribing physician indicated on the form. Thus, although the forms probably have a role in guiding physicians, time pressures and other factors lead to unsafe prescribing practices. Because the form was used for almost all patients, we are unable to determine whether the risk of adverse events decreased with its use.

We support the idea of improving communication between multidisciplinary members of the health care team. MacLean highlights the changes in medication regimens that are often made during and after a hospital stay. Frequently, this information is not communicated effectively to patients,² pharmacists or community physicians.³ The need to reconcile medication regimens before and after the hospital stay and the need to improve communications pertaining to medication use are obvious. However, translating these needs into practical, effective solutions will require substantially more investment than changes in paper forms. Although unproven, it is possible that bet-

ter hospital information systems will be required, e.g., through computerized physician order entry⁴ or automated discharge summary generation.⁵

Alan J. Forster

Carl van Walraven

The Ottawa Hospital
The Ottawa Health Research Institute
Ottawa, Ont.

References

1. Forster AJ, Clark HD, Menard A, Dupuis N, Chernish R, Chandok N, et al. Adverse events among medical patients after discharge from hospital. *CMAJ* 2004;170(3):345-9.
2. Forster AJ, Bates DW, Gandhi T, Murff HJ, Peterson J. Adverse drug events following hospital discharge [abstract]. *J Gen Intern Med* 2003;18(Suppl 1):282.
3. van Walraven C, Weinberg AL. Quality assessment of a discharge summary system. *CMAJ* 1995;152(9):1437-42.
4. Bates DW, Leape LL, Cullen DJ, Laird N, Petersen LA, Teich JM, et al. Effect of computerized physician order entry and a team intervention on prevention of serious medication errors. *JAMA* 1998;280(15):1311-6.
5. van Walraven C, Laupacis A, Seth R, Wells G. Dictated versus database-generated discharge summaries: a randomized clinical trial. *CMAJ* 1999;160(3):319-26.

DOI:10.1053/cmaj.1040686

Correction

A recent recommendation statement on postmenopausal hormone replacement therapy for primary prevention of cardiovascular and cerebrovascular disease¹ should have included the following byline: "Beth L. Abramson and the Canadian Task Force on Preventive Health Care." Beth Abramson is Assistant Professor of Medicine, University of Toronto, and Director of the Cardiac Prevention Centre and Women's Cardiovascular Health, St. Michael's Hospital, Toronto, Ont.

Reference

1. Abramson BL, Canadian Task Force on Preventive Health Care. Postmenopausal hormone replacement therapy for primary prevention of cardiovascular and cerebrovascular disease: recommendation statement from the Canadian Task Force on Preventive Health Care. *CMAJ* 2004;170(9):1388-9.

New letters submission process

CMAJ's enhanced eLetters feature is now the portal for all submissions to our letters column. To prepare an eLetter, visit www.cmaj.ca and click "Submit a response to this article" in the box near the top right-hand corner of any *eCMAJ* article. All eLetters will be considered for publication in the print journal.

Letters written in response to an article published in *CMAJ* are more likely to be accepted for print publication if they are submitted within 2 months of the article's publication date. Letters accepted for print publication are edited for length (usually 250 words) and house style.