ing a benefit over placebo is analogous to not treating meningitis in pregnancy because there are no randomized trials to demonstrate that penicillin is better than placebo in pregnancy.

Zitner and Bischoff quote one of us as having said, elsewhere, that "What we found was that [among] pregnant women who use Paxil through pregnancy until birth, their offspring are more likely to have several stormy weeks at infancy." This was taken out of context, as they omitted the accompanying statement that "Many of these babies have to stay in the hospital for two to three weeks after they're born, but they suffer no long-term health effects."⁵

Although it may be possible for some women to avoid taking antidepressants as Zitner and Bischoff suggest, they do not offer an alternative approach for the substantial number of women who have major depressive symptoms during pregnancy. Antidepressants continue to be prescribed and it is important that pregnant women and their health care providers have accurate information upon which to base an informed decision regarding therapy.

Gideon Koren

Motherisk Program
Hospital for Sick Children
Toronto, Ont.

Doreen Matsui

Department of Pediatrics
University of Western Ontario
London, Ont.

Adrienne Einarson

Motherisk Program
Hospital for Sick Children
Toronto, Ont.

David Knopper

Department of Pediatrics
University of Western Ontario
London, Ont.

Meir Steiner

Department of Psychiatry
and Behavioural Neurosciences
McMaster University
Hamilton, Ont.

REFERENCES

1. Koren G, Matsui D, Einarson A, et al. Is maternal use of selective serotonin reuptake inhibitors in the third trimester of pregnancy harmful to neonates? *CMAJ* 2005;172(11):1457-9.
2. Huot RL, Brennan PA, Stowe ZN, et al. Negative effect in offspring of depressed mothers is predicted

by infant cortisol levels at 6 months and maternal depression during pregnancy, but not postpartum. *Ann NY Acad Sci* 2004;1032:234-6.

3. Nordeng H, Spigset O. Treatment with selective serotonin reuptake inhibitors in the third trimester of pregnancy: effects on the infant. *Drug Saf* 2005; 28:565-81.
4. Ruchkin V, Martin A. SSRIs and the developing brain. *Lancet* 2005;365 (9458):451-3.
5. Paxil in late pregnancy may cause problems. Available: www.healthplace.com/communities/depression/treatment/antidepressants/articles/004.asp (accessed 2005 Oct 12).

DOI:10.1503/cmaj.1050171

A positive prognosis

We read with interest the recent report by Krishna Sharma¹ of a case of malignant fibrous histiocytoma, metastatic to the lung, with spontaneous expectoration of large tumour fragments. We felt it would be illustrative to present a similar case with a much more favourable outcome.

This patient presented in 1975 at the age of 42 years with a slowly enlarging mass, involving the right patellar tendon. A biopsy revealed it to be a malignant sarcoma, and a subsequent wide local excision was accomplished with clear margins. Pathology review confirmed the diagnosis of malignant fibrous histiocytoma. No adjuvant treatment was administered.

Five years after primary resection, the patient developed pulmonary nodules, one adjacent to each hilum, visible on both routine posteroanterior chest radiographs and lung tomograms. He was asymptomatic, and no further treatment was recommended. Over the next 8 months, he developed mild but progressive wheezing. This culminated in the spontaneous expectoration of a tumour nodule described as a plug of tissue measuring 2.5 × 1 cm and found to be histologically identical to his primary cancer. The wheezing completely resolved at this point but because of the enlargement of his remaining disease site, with the largest nodule measuring 5 cm in diameter, a course of radiation therapy was recommended. He received 30 Gy in 15 fractions, encompassing both hila and the carina, using cobalt 60. This was well tolerated aside from transient fatigue. The patient's tumour masses began to shrink promptly, and he was eventually left with a small amount of residual scarring near his left hilum.

When last seen in follow-up in 2003, some 22 years after the spontaneous expectoration of one of his lung nodules and subsequent "palliative" radiation treatment of his residual disease, the patient remained alive and well, without disease recurrence.

Anthony Brade

Brian O'Sullivan

Bernard Cummings

Department of Radiation Oncology

Ian Quirt

Department of Medical Oncology

Princess Margaret Hospital

University Health Network

University of Toronto

Toronto, Ont.

REFERENCE

1. Sharma KB. Spontaneous expectoration of tumour mass. *CMAJ* 2005;172(9):1182.

DOI:10.1503/cmaj.1050141

Predicting cardiac outcomes

Despite substantial advances in the diagnosis of suspected acute coronary syndromes, significant challenges persist.¹ Andrew Worster and colleagues recently reported in *CMAJ* that ischemia-modified albumin (IMA) was a poor predictor of cardiac outcomes in patients with potential cardiac ischemia symptoms.² The authors tested 2 thresholds for IMA: 85 µ/mL, as suggested by the manufacturer, and 80 µ/mL. It is important to point out, however, that IMA levels vary considerably, even among healthy individuals. Taking these variations into account may improve the predictive characteristics of IMA.

We examined 35 healthy men (age range 25–54 years) recruited from the general public who had not had a myocardial infarction. Using standard laboratory techniques we found that the average resting IMA concentration was 94 µ/mL (97.5% confidence interval 84–104 µ/mL). IMA concentration was significantly and inversely correlated with serum albumin but not with creatinine concentration. Serum IMA concentration was also significantly associated with serum lactate concentration.³ Taking these and other known factors into