has developed a set of guidelines for industry-sponsored clinical trial and epidemiology contract research in Canada in collaboration with 6 multinational vaccine manufacturers.⁵ These guidelines describe the roles of the academic and public health investigator in protocol development, access to data, data management, data analysis, creation of the study report and publication in peer-reviewed journals. The guidelines conform to the recommendations of the ICMJE and aim to ensure that the academic or public health investigators have full and unrestricted access to data and play a leading role in the publication of study results. We hope that these guidelines will address the deficiencies documented in recent evaluations of clinical trial agreements and will serve as a model for the interaction between academia, public health and the pharmaceutical industry.

Scott A. Halperin
David Scheifele
Bernard Duval
Brian Ward

On behalf of the Canadian Association
 for Immunization Research
 and Evaluation (CAIRE)
 Vancouver, BC

REFERENCES

1. Davidoff F, DeAngelis CD, Drazen JM, et al. Sponsorship, authorship and accountability. *CMAJ* 2001;165(6):786-8.
2. Schulman KA, Seils DM, Timbie JW, et al. A national survey of provisions in clinical-trial agreements between medical schools and industry sponsors. *N Engl J Med* 2002;347:1335-41.
3. Mello MM, Clarridge BR, Studdert DM. Academic medical centers' standards for clinical-trial agreements with industry. *N Engl J Med* 2005;352:2202-10.
4. Steinbrook R. Gag clauses in clinical-trial agreements. *N Engl J Med* 2005;352:2160-2.
5. Halperin SA, Scheifele D, Duval B, et al. Canadian Association for Immunization Research and Evaluation (CAIRE) Guidelines for Industry Sponsored Clinical Trial and Epidemiology Contract Research. *Hum Vaccines* 2005;1(4):140-2.

DOI:10.1503/cmaj.1050172

Refresher on rubella

The authors of a recent public health column in *CMAJ* state that "women of childbearing age should be given rubella vaccine unless they have proof of immunity, and they should be advised to avoid pregnancy for 3 months after vaccination."¹ However, a public

health document cited in the column states that "women of childbearing age should be advised to avoid pregnancy for 1 month after immunization. This recommendation is based on the duration of viremia after natural infection and evidence of vaccine safety."

I recently saw a patient who tested negative for rubella antibodies at her first visit. When recalled for a measles, mumps and rubella vaccination, she told me that she wanted to have a child and did not want to wait too long before trying to conceive. I thought a 3-month wait was required but found the 1-month recommendation² when I double-checked the requirement.

Michelle Greiver
 Family Physician
 Toronto, Ont.

REFERENCES

1. Weir E, Sider D. A refresher on rubella. *CMAJ* 2005;172(13):1680-1.
2. Vaccine preventable diseases — rubella. Ottawa: Immunization and Respiratory Infections Division, Public Health Agency; 2002. Available: www.phac-aspc.gc.ca/dird-dimr/vpd-mev/rubella_e.html (accessed 2005 Oct 14).

DOI:10.1503/cmaj.1050156

[The author responds:]

I thank Michelle Greiver for her attention to our column on rubella.¹ In the first draft of this article Doug Sider and I quoted the Public Health Agency of Canada's advice² (which appeared to be based on information in the 2002 Canadian Immunization Guide³) that women should avoid pregnancy for 1 month after receiving the rubella vaccine. However, a peer reviewer challenged this statement because one of the bibles on communicable disease, the *Control of Communicable Diseases Manual*,⁴ states that after immunization for rubella women should prevent pregnancy for 3 months. In the end we decided to relay the more cautious message.

When I try to assess and communicate risk, I usually ask 3 questions: Is there a hazard? If so, what is the magnitude of the adverse outcome? What is the likelihood that the adverse outcome will occur? In this case, there is a hazard associated with immunizing a

pregnant woman; the magnitude of the adverse outcome is potentially very great (congenital rubella syndrome) but the likelihood that it will occur is extremely low (bordering on theoretical). Because the potential adverse outcome is so serious we decided to relay the more conservative recommendation. However, given the extremely low likelihood that it will occur, one could as easily argue that advising women to avoid pregnancy for 1 month after immunization is appropriate.

Erica Weir
CMAJ

REFERENCES

1. Weir E, Sider D. A refresher on rubella. *CMAJ* 2005;172(13):1680-1.
2. Vaccine preventable diseases — rubella. Ottawa: Immunization and Respiratory Infections Division, Public Health Agency; 2002. Available: www.phac-aspc.gc.ca/dird-dimr/vpd-mev/rubella_e.html (accessed 2005 Oct 14).
3. National Advisory Committee on Immunization. *Canadian immunization guide*. 6th ed. Ottawa: Health Canada; 2002. Available: www.phac-aspc.gc.ca/publicat/cig-gci (accessed 2005 Oct 14).
4. Chin J, editor. *Control of communicable diseases manual: an official report of the American Public Health Association*. 17th ed. Washington (DC): American Public Health Association; 2000.

DOI:10.1503/cmaj.1050157

Vitamin B₁₂ and homocysteine

In their analysis of the relationship between carotid plaque area and vitamin B₁₂ status,¹ Julie Robertson and colleagues define vitamin B₁₂ deficiency as a serum concentration of less than 258 pmol/L, with a plasma homocysteine concentration of 14 µmol/L or more or a plasma methylmalonic acid (MMA) level of 271 nmol/L or more. We question the use of these thresholds and the conclusions that are based on them.

Their threshold for vitamin B₁₂ deficiency of 258 pmol/L is almost the same as the sample median value of 253 pmol/L. This value is exceedingly high and probably inappropriately labeled many study patients as having a vitamin B₁₂ deficiency. Among 11 000 elderly women in Ontario and British Columbia, the mean and fifth centile serum vitamin B₁₂ concentrations were 300 and 118 pmol/L, respectively, after

folic acid food fortification.² Similar values were found among Ontario women of reproductive age.³

We previously showed that plasma total homocysteine levels can be relatively insensitive for detecting vitamin B₁₂ deficiency.⁴ A homocysteine concentration at or above 15 µmol/L has positive and negative predictive values of 7.4% and 97.2%, respectively, for the detection of a serum vitamin B₁₂ concentration below 120 pmol/L. For vitamin B₁₂ levels between 120 and 150 pmol/L, the corresponding predictive values are just 6.3% and 94.0%.⁴ In Fig. 1 of the article by Robertson and colleagues the mean homocysteine concentration was just 12 µmol/L even in the quartile of patients with the lowest serum vitamin B₁₂ concentrations (below 203 pmol/L). Thus, it is unlikely that vitamin B₁₂ deficiency could be efficiently detected on the basis of a homocysteine measurement equal to or greater than 15 µmol/L.

Their lack of use of a comprehensive and suitable definition of vitamin B₁₂ leads us to question the authors' conclusions that "vitamin B₁₂ deficiency is surprisingly common among patients with vascular disease" and that "low serum vitamin B₁₂ levels are a major determinant of elevated homocysteine levels and increased carotid plaque area."¹ With nearly 50% of patients omitted from their major analysis, could not age and serum creatinine alone have explained some of their findings?

Joel G. Ray

Assistant Professor
Departments of Medicine, Health
Policy Management and Evaluation,
and Obstetrics and Gynecology

David E.C. Cole

Professor
Department of Clinical Pathology
University of Toronto
Toronto, Ont.

REFERENCES

1. Robertson J, Iemolo F, Stabler SP, et al. Vitamin B₁₂, homocysteine and carotid plaque in the era of folic acid fortification of enriched cereal grain products. *CMAJ* 2005;172(12):1569-73.
2. Ray JG, Vermeulen MJ, Langman LJ, et al. Persis-

tence of vitamin B₁₂ insufficiency among elderly women after folic acid food fortification. *Clin Biochem* 2003;36:387-91.

3. Ray JG, Vermeulen MJ, Boss SC, et al. Increased red cell folate concentrations in women of reproductive age after Canadian folic acid food fortification. *Epidemiology* 2002;13:238-40.
4. Ray JG, Cole DE, Boss SC. An Ontario-wide study of vitamin B₁₂, serum folate, and red cell folate levels in relation to plasma homocysteine: Is a preventable public health issue on the rise? *Clin Biochem* 2000;33:337-43.

DOI:10.1503/cmaj.1050153

In examining the relationship of serum vitamin B₁₂ levels to homocysteine levels,¹ Julie Robertson and colleagues highlight the utility of homocysteine in the clinical investigation of vitamin B₁₂ deficiency.

Both methylmalonic acid and homocysteine are elevated in vitamin B₁₂ deficiency. The measurement of methylmalonic acid requires not readily available methods involving mass spectrometry, whereas homocysteine is readily measured in the clinical laboratory by fluorescent polarization, enzymatic assays and high-pressure liquid chromatography. There is a strong correlation between methylmalonic acid and homocysteine levels, and the sensitivity of homocysteine levels for identifying vitamin B₁₂ deficiency is greater than 95%.²⁻⁴ Measurement of creatinine, folate and vitamin B₁₂ levels and the patient's response to treatment will clarify the cause of the elevated homocysteine levels. At my institution we no longer routinely offer methylmalonic acid measurement as the diagnostic efficiencies of methylmalonic acid and homocysteine seem similar for the identification of a functional vitamin B₁₂ deficiency.

Robertson and associates state that 17.3% of their population (73/421 cases) had a vitamin B₁₂ deficiency. Of the 50 patients in whom methylmalonic acid was measured, 59% had elevated methylmalonic acid levels, confirming a functional vitamin B₁₂ deficiency. It would have been interesting to know what the homocysteine levels were in these 73 subjects and to conduct a receiver-operator curve analysis to determine the efficiency of methyl-

malonic acid and homocysteine for identifying vitamin B₁₂ deficiency. If all the patients with elevated methylmalonic acid levels also had elevated homocysteine levels, this would support the idea that homocysteine alone (without methylmalonic acid) can be used to assess vitamin B₁₂ deficiency and its response to treatment.

Brian M. Gilfix

Associate Physician
Division of Clinical Biochemistry
Royal Victoria Hospital
Montréal, Que.

REFERENCES

1. Robertson J, Iemolo F, Stabler SP, et al. Vitamin B₁₂, homocysteine and carotid plaque in the era of folic acid fortification of enriched cereal grain products. *CMAJ* 2005;172(12):1569-73.
2. Holleland G, Schneede J, Ueland PM, et al. Cobalamin deficiency in general practice. Assessment of the diagnostic utility and cost-benefit analysis of methylmalonic acid determinations in relation to current diagnostic strategies. *Clin Chem* 1999;45(2):189-98.
3. Refsum H, Smith AD, Ueland PM, et al. Facts and recommendations about total plasma homocysteine determinations: an expert opinion. *Clin Chem* 2004;50(1):3-32.
4. Savage DG, Lindebaum J, Stabler SP, et al. Sensitivity of serum methylmalonic acid and total homocysteine determinations for diagnosing cobalamin deficiencies. *Am J Med* 1994;96:239-46.

DOI:10.1503/cmaj.1050170

Correction

A news article¹ concerning the federal government's decision not to renew funding to the Canadian Network for Vaccines and Immunotherapeutics stated that the network's early success included the "beginning of the first Canadian clinical trial for a therapeutic HIV vaccine." Although the final results of this research were presented at a conference organized by the network, the research itself was sponsored by the Canadian HIV Trials Network.

REFERENCE

1. Eggertson L. Vaccine network surprised by funding cut. *CMAJ* 2005;173(7):741-2.

DOI:10.1503/cmaj.051391