practice of medicine, and the SOGC finds it important to disseminate this information to our members. Better for health professionals to receive medical updates from a specialty society rather than directly from industry itself or, worse, direct-to-consumer advertising from industry. The article was independently produced with no input or review from industry. Its dissemination was provided by an unrestricted educational grant. The fact that this article sheds a positive light on a new product does not indicate a conflict of interest. What we at the SOGC advocate for is up-to-date accurate dissemination of information to our members, and this document accomplishes this.

Vyta Senikas

Associate Executive Vice-President
Society of Obstetricians and
Gynaecologists of Canada
Ottawa, Ont.

Competing interests: None declared.

DOI:10.1503/cmaj.1050158

Well-water maintenance

We congratulate Erica Weir for having recently drawn attention to well-water maintenance.¹ We have some additional information to help physicians interpret the results of reports on well-water quality.

Although we agree with the list of potential contaminants of well water presented by Weir in Box 1, it is worth mentioning that routine monitoring of water microbiological quality entails the detection of the nonpathogenic coliforms, the total and thermotolerant (fecal) coliforms. Nonpathogenic *Escherichia coli* are the most common coliform in human and animal feces and they are recognized as the best index of recent fecal contamination of surface water and groundwater.^{2,3} In the presence of thermotolerant (fecal) coliforms or *E. coli*, a boil-water advisory is advisable until the water is treated and disinfected.

Total coliforms, which might be present in the general environment (in soil and plants) without fecal contamination, are usually considered as an indicator of the vulnerability of groundwa-

ter to microbiological contamination. If total coliforms are detected, a boil-water advisory is usually not recommended but inspection of the well and more frequent analysis of the water for *E. coli* is advisable.

Nonpathogenic intestinal enterococci (a subgroup of fecal streptococci) appear to survive longer in the environment than *E. coli* and are being proposed as microbiological indicators of groundwater quality. They are also considered to be an index of fecal contamination,^{4,5} although they are not totally specific to animal or human feces.

Pathogens are rarely measured in drinking water because they are expensive to detect and detection methods have not yet been standardized.⁶ Nevertheless, pathogen testing is extremely useful for outbreak investigation.⁷

Patrick Levallois

Direction des risques biologiques,
environnementaux et occupationnels
Institut national de santé publique
du Québec
Sainte-Foy, Que.

Pierre Payment

Institut Armand-Frappier
Institut national de la recherche
scientifique
Laval, Que.

REFERENCES

1. Weir E. Well-water maintenance. *CMAJ* 2005;172(11):1438.
2. Federal-Provincial-Territorial Committee on Drinking Water. *Bacteriological quality. guidelines for Canadian drinking water quality: supporting documentation*. Ottawa: Health Canada; 2002. Available: www.hc-sc.gc.ca/hecs-sesc/water/publications/bacteriological_quality/toc.htm (accessed 2005 Jul 7).
3. Ministère du développement durable, environnement et parcs. *La qualité de l'eau de mon puits*. Québec: Gouvernement du Québec; 2002. www.menv.gouv.qc.ca/eau/potable/depliant/index.htm#qualite (accessed 2005 Jul 7).
4. World Health Organization. *WHO guidelines for drinking water quality*. 3rd ed. Geneva: The Organization; 2004. Available: www.who.int/water_sanitation_health/dwq/guidelines/en/index.html (accessed 2005 Jul 7).
5. Groupe scientifique sur l'eau. *Entérocoques et streptocoques fécaux*. Sainte-Foy (Que): Institut national de santé publique du Québec; 2002. Available: www.inspq.qc.ca/pdf/publications/198-CartableEau/EnteroCoques.pdf (accessed 2005 Jul 7).
6. Medema GJ, Payment P, Dufour A, Robertson W, Waite M, Hunter P, et al. Safe drinking water: an ongoing challenge. In: Dufour A, Snozzi M, Köster W, Bartram J, Ronchi E, Fewtrell L, editors. *Assessing microbial safety of drinking water: improving approaches and methods. WHO drinking water quality series*. London: IWA Publishing; 2003. p. 11-46.
7. De Serres G, Cromeans TL, Levesque B, Brassard

N, Barthe C, Dionne M, et al. Molecular confirmation of hepatitis A virus from well water: epidemiology and public health implications. *J Infect Dis* 1999;179:37-43.

DOI:10.1503/cmaj.1050148

HIV/AIDS and leishmaniasis coinfection in Ethiopia

Aranka Anema and Koert Ritmeijer¹ describe their practical and effective strategy for treating visceral leishmaniasis (VL) in patients coinfecting with HIV/AIDS in Ethiopia. Relying on low serologic titres of $\leq 1:400$ to exclude VL is problematic, because titres of $\leq 1:400$ might well be the result of patients with leishmaniasis having a grossly inadequate serologic response.

Coinfection with HIV and VL is documented to be associated with a poor serologic response. In Spain, among 120 patients with VL, including 80 coinfecting with HIV, the serologic response was significantly lower among those coinfecting with HIV.² An identical scenario in Kafta Humera Woreda would lead to an underdiagnosis of VL and spoil the utility of an efficient decentralized diagnostic and therapeutic service.

It would be worthwhile to investigate at least some of those with serological titres of $\leq 1:400$ for *Leishmania donovani* in their tissue aspirates. If costs are prohibitive in Ethiopia, then maybe the international community could help with this assay.

Subhash C. Arya

Nirmala Agarwal
Sant Parmanand Hospital
Delhi, India

REFERENCES

1. Anema A, Ritmeijer K. Treating HIV/AIDS and leishmaniasis coinfection in Ethiopia. *CMAJ* 2005;172(11):1434-5.
2. Pintado V, Martín-Rabadán P, Rivera ML, Moeno S, Bouza E. Visceral leishmaniasis in human immunodeficiency virus (HIV)-infected and noninfected patients. A comparative study. *Medicine* 2001;80:54-73.

DOI:10.1503/cmaj.1050140

The direct agglutination test (DAT) is not the only diagnostic test used

in the program we described.¹ Even if a case of VL and HIV coinfection was missed by DAT, the diagnostic algorithm used in Ethiopia would still identify it, because in the case of a negative DAT result, but persisting signs and symptoms of VL, a splenic aspiration is indicated. DAT is used as a first-line test to reduce the number of tissue aspirates required for screening clinically suspicious cases.

The results from European studies cannot be extrapolated to African VL, because the patient characteristics are quite different. For example, in the study by Pintado and colleagues,² which Subhash Arya and Nirmala Agarwal refer to, patients were mainly injection drug users who were profoundly immunocompromised. An evaluation of DAT in Ethiopian patients with VL (with and without HIV infection) concluded that in contrast to the observations made in Europe, DAT in Ethiopia remains reasonably sensitive in diagnosing VL during HIV coinfection.³ These findings are confirmed by unpublished data from the Médecins Sans Frontières miltefosine study in Kafta Humera Woreda, which show that DAT titres in cases of HIV coinfection did not differ from those in cases where there was no coinfection, that is, there was no shift in the mean and distribution of DAT titres.

Aranka Anema
Médecins Sans Frontières
Canada
Koert Ritmeijer
Médecins Sans Frontières
Holland

REFERENCES

1. Anema A, Ritmeijer K. Treating HIV/AIDS and leishmaniasis coinfection in Ethiopia. *CMAJ* 2005; 172(11):1434-5.
2. Pintado V, Martin-Rabadou P, Rivera ML, Moeno S, Bouza E. Visceral leishmaniasis in human immunodeficiency virus (HIV)-infected and noninfected patients. A comparative study. *Medicine* 2001;80:54-73.
3. Hailu A, Berhe N. The performance of direct agglutination tests (DAT) in the diagnosis of visceral leishmaniasis among Ethiopian patients with HIV co-infection. *Ann Trop Med Parasitol* 2002;96(1):25-30.

DOI:10.1503/cmaj.1050167

Are FASD guidelines practical and sustainable?

Christine Loock and associates¹ recommend that children with suspected fetal alcohol spectrum disorder (FASD) undergo a "comprehensive, multidisciplinary assessment." The assessment outlined in recommendation 3.1 of the supplement presenting Canadian guidelines on diagnosis of FASD² is comprehensive but is neither justified by evidence nor sustainable within existing resources. If the prevalence of this disorder is around 1%, as suggested,² more than 400 such children per year would need assessment in Alberta alone.

On the basis of data in the 2004 annual reports of the Alberta Children's Hospital and the Glenrose Rehabilitation Hospital, and allowing for the fact that children are also seen in other regional assessment centres, I estimate that approximately 800 preschool children are now seen annually by our tertiary developmental assessment centres, so this would add another 50% to their workload. A small number of children with FASD are assessed by our local tertiary-level team each year, but the guideline implies that children currently being seen by family doctors, general pediatricians or subspecialists (e.g., developmental pediatricians, clinical geneticists, child psychiatrists), in conjunction with early intervention workers, school psychologists, community speech-language pathologists and others, are receiving inadequate diagnosis. This is a very important issue, given that our provincial education ministry provides extra funding to school districts for children with a diagnosis of FASD; one risk of these guidelines is that none of these children will meet the diagnostic criteria for funding.

The "tertiary bias" of the guideline is further illustrated by recommendation 1.4: among the "appropriate professionals" not mentioned (for children with other developmental disabilities) are colleagues from family medicine or general pediatrics. Even subspecialists such as myself often function in a community-based role. While access to the

full multidisciplinary team is essential for some of the children I see, I believe that the diagnostic assessment I provide is quite appropriate for many children within a more limited scope (i.e., working collaboratively with community practitioners).

I suggest that children with a confirmed history of exposure to alcohol and relevant developmental, behavioural and phenotypic findings should receive a diagnosis within the spectrum of FASD by anyone competent to do so, including an experienced primary care physician. Children for whom there is genuine doubt about the diagnosis should be referred for further assessment by an appropriate multidisciplinary team.

Recommendations for more complex diagnostic processes require clear evidence concerning cost and benefit, sensitivity and specificity, and validity and reliability.

Keith J. Goulden
Neurodevelopmental Pediatrics
Glenrose Rehabilitation Hospital
Edmonton, Alta.

REFERENCES

1. Loock C, Conry J, Cook JL, Chudley AE, Rosales T. Identifying fetal alcohol spectrum disorder in primary care. *CMAJ* 2005;172(5):628-30.
2. Chudley AE, Conry J, Cook JL, Loock C, Rosales T, LeBlanc N. Fetal alcohol spectrum disorder: Canadian guidelines for diagnosis. *CMAJ* 2005;172(5 Suppl):S1-21.

DOI:10.1503/cmaj.1050107

[The authors respond:]

The FASD guidelines¹ and accompanying article² should not be interpreted as implying a "tertiary bias," as suggested by Keith Goulden. On the contrary, trained and functional FASD diagnostic teams have recently been established in smaller communities throughout Canada and are providing excellent service outside of tertiary care facilities, often without a subspecialist's direct involvement.

According to the diagnostic process outlined in the guidelines, any physician "specifically trained in FASD diagnosis" can be a member of the team. Such multidisciplinary di-