nutrition standards and most effective approaches to front-of-pack labelling.

Bill Jeffery LLB

National Coordinator, Centre for Science in the Public Interest (Canada), Ottawa, Ont.

Competing interests: None declared.

REFERENCES

1. Truscott A. Checking up on Health Check. *CMAJ* 2008;178:386-7.
2. Katz, DL. *FAQs about the ONQI*. Derby (CT): Yale Griffin Prevention Clearance Center; 2007. Available: www.griffinhealth.org/Research/ONQIpages/FAQs.aspx (accessed 2007 Mar 19).
3. Arambepola C, Scarborough P, Rayner M. Validating a nutrient profile model. *Public Health Nutr* 2008;11:371-8.
4. Heart and Stroke Foundation of Canada. *Check for Health Check. Frequently asked questions*. Ottawa: The Foundation; 2008. Available: www.healthcheck.org/en/about-health-check/frequently-asked-questions.html (accessed 2008 Mar 19).

DOI:10.1503/cmaj.1080029

[Dr. Freedhoff responds:]

The Heart and Stroke Foundation of Canada's Health Check program was recently examined in a news piece in *CMAJ*.¹ Although it is true that the nutrient criteria for this program have recently been revised,² they still permit products to contain tremendous amounts of sodium and sugar.

Health Check's criteria currently allow individual products to contain between 21% and 57% of the maximal daily sodium intake recommended by Health Canada and between 32% and 87% of that recommended by Blood Pressure Canada.³ These allowances reflect tremendous hypocrisy, especially given that in 2007 the Heart and Stroke Foundation, along with Blood Pressure Canada, the Canadian Hypertension Education Program, the Canadian Hypertension Society and the Société Québécoise d'Hypertension Artérielle, urged Canadians to avoid foods containing more than 10% of the maximal daily recommended amount of sodium.⁴

Using a diet made up entirely of items endorsed by the Health Check program, I found that the program

allows people to easily consume more than 4000 mg of sodium and 190 g of sugar daily, with sugar accounting for over 30% of total calories (calculations available at www.cmaj.ca/cgi/eleletters/178/4/386#18544). These values are nearly double the maximal sodium intake recommended by Health Canada, nearly triple the maximal sodium intake recommended by Blood Pressure Canada and triple the World Health Organization's recommended maximal daily sugar allowance.⁵

It would be tragic if we were blinded by the Heart and Stroke Foundation's reputation and adopted Health Check as Canada's national front-of-package labelling program, because Health Check's minimalist and lax nutritional criteria sorely fail to guide Canadians to healthier choices. It is time for Canadians to demand that national policies and recommendations be grounded in evidence.

Yoni Freedhoff MD

Medical Director, Bariatric Medical Institute, Ottawa, Ont.

Competing interests: None declared.

REFERENCES

1. Truscott A. Checking up on Health Check. *CMAJ* 2008;178:386-7.
2. Heart and Stroke Foundation of Canada. *Check for Health Check. Nutrient criteria*. Available: www.healthcheck.org/en/nutritional-information/nutrient-criteria-grocery.html (accessed 2008 Mar 14).
3. Blood Pressure Canada. *Policy: sodium*. Available: <http://hypertension.ca/bpc/wp-content/uploads/2007/10/bpc-sodium-policy-with-endorsements-clean.pdf> (accessed 2008 Mar 14).
4. Blood Pressure Canada, Heart and Stroke Foundation of Canada, Canadian Hypertension Education

Program, et al. *Hypertension: 2007 public recommendations*. Available: <http://hypertension.ca/bpc/wp-content/uploads/2007/11/chep2007march-locked.pdf> (accessed 2008 Mar 14).

5. World Health Organization. *Diet, nutrition and the prevention of chronic diseases: report of a joint WHO/FAO expert consultation*. WHO Tech Rep Ser no 916. Geneva: The Organization; 2003.

DOI:10.1503/cmaj.1080026

Corrections

An article in The Left Atrium in the Feb. 26 issue about Bolivian travelling doctors contained an error and an omission.¹ The correct spelling of the term for these individuals is *medicos viajeros Bolivianos*. In addition, the coauthor of this article was Andrew G. Malleson MBBS, Department of Psychiatry (retired), University Health Network, University of Toronto, Toronto, Ont. *CMAJ* apologizes for this error and any inconvenience it may have caused.

REFERENCE

1. Stewart D. *Medicos viajeros Bolivianos*. *CMAJ* 2008;178:602.

DOI:10.1503/cmaj.080338

A Dispatch from the Medical Front in the March 11 issue should have identified Tamale as being located in Ghana. Because of an editing error, it was identified as being in Uganda.¹

REFERENCE

1. Lin, Daren. Soup truck. *CMAJ* 2008;178:673.

DOI:10.1503/cmaj.080518

Letters submission process

CMAJ's enhanced letters feature is now the portal for all submissions to our letters column. To prepare a letter, visit www.cmaj.ca and click "Submit a response to this article" in the box near the top right-hand corner of any *CMAJ* article. All letters will be considered for publication in the print journal.

Letters written in response to an article published in *CMAJ* are more likely to be accepted for print publication if they are submitted within 2 months of the article's publication date. Letters accepted for print publication are edited for length (usually 250 words) and house style.