

Arthroscopic surgery for degenerative tears of the meniscus: a systematic review and meta-analysis

Moin Khan MD, Nathan Evaniew MD, Asheesh Bedi MD, Olufemi R. Ayeni MD MSc, Mohit Bhandari MD PhD

ABSTRACT

Background: Arthroscopic surgery for degenerative meniscal tears is a commonly performed procedure, yet the role of conservative treatment for these patients is unclear. This systematic review and meta-analysis evaluates the efficacy of arthroscopic meniscal débridement in patients with knee pain in the setting of mild or no concurrent osteoarthritis of the knee in comparison with nonoperative or sham treatments.

Methods: We searched MEDLINE, Embase and the Cochrane databases for randomized controlled trials (RCTs) published from 1946 to Jan. 20, 2014. Two reviewers independently screened all titles and abstracts for eligibility. We assessed risk of bias for all included studies and pooled outcomes using a random-effects model. Outcomes (i.e., function and pain relief) were dichotomized to short-term (< 6 mo) and long-term (< 2 yr) data.

Results: Seven RCTs ($n = 805$ patients) were included in this review. The pooled treatment

effect of arthroscopic surgery did not show a significant or minimally important difference (MID) between treatment arms for long-term functional outcomes (standardized mean difference [SMD] 0.07, 95% confidence interval [CI] -0.10 to 0.23). Short-term functional outcomes between groups were significant but did not exceed the threshold for MID (SMD 0.25, 95% CI 0.02 to 0.48). Arthroscopic surgery did not result in a significant improvement in pain scores in the short term (mean difference [MD] 0.20, 95% CI -0.67 to 0.26) or in the long term (MD -0.06, 95% CI -0.28 to 0.15). Statistical heterogeneity was low to moderate for the outcomes.

Interpretation: There is moderate evidence to suggest that there is no benefit to arthroscopic meniscal débridement for degenerative meniscal tears in comparison with nonoperative or sham treatments in middle-aged patients with mild or no concomitant osteoarthritis. A trial of nonoperative management should be the first-line treatment for such patients.

Competing interests:

Mohit Bhandari declares consultancy payments from Smith & Nephew, Stryker, Amgen, Zimmer, Moximed and Bioventus, and grant support from Smith & Nephew, DePuy, Eli Lilly and Bioventus. No other competing interests were declared.

This article has been peer reviewed.

Correspondence to:

Moin Khan, moinkhanmd@gmail.com

CMAJ 2014. DOI:10.1503/cmaj.140433

Arthroscopic meniscal débridement is one of the most commonly performed procedures in orthopedic surgery. More than 700 000 such procedures are performed each year in the United States, and more than 4 million are performed each year worldwide, with substantial economic and social burdens.¹⁻⁶ Many patients who undergo arthroscopic meniscal débridement have concurrent osteoarthritis, and orthopedic surgeons are often challenged to determine the true cause of patients' symptoms: the meniscal tear, osteoarthritis or a combination of both.⁷

Although 2 well-designed randomized controlled trials (RCTs)^{8,9} have shown a lack of efficacy for arthroscopic surgery in patients with severe and advanced knee arthritis, many patients present with degenerative meniscal tears and mild or minimal concurrent osteoarthritis.¹⁰ Patients with degenerative meniscal tears in the setting of mild osteoarthritis may experience functional improvement or pain relief with

arthroscopic surgery,¹¹⁻¹⁴ but the role of conservative treatment is unclear.¹⁵⁻¹⁷ Arthroscopic surgery involves the potential for complications, which must be weighed against the prognosis for relief from presenting symptoms.^{6,18}

The objective of this systematic review and meta-analysis was to evaluate the efficacy of arthroscopic meniscal débridement in comparison with nonoperative or sham treatments in patients with degenerative meniscal tears and knee pain with regard to function and pain relief in the short term (< 6 mo) and long term (< 2 yr).

Methods

We conducted this study according to the methods of the *Cochrane Handbook for Systematic Reviews of Interventions*.¹⁹ The findings are reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement.²⁰

Eligibility criteria

We included RCTs that 1) involved patients of any age or sex with degenerative meniscal tears and mild or no concurrent osteoarthritis presenting with knee pain, and 2) compared arthroscopic meniscal débridement (with or without concurrent articular débridement) with nonoperative treatments.

No restriction was made regarding publication date, language, presence or absence of co-interventions, specific nonoperative intervention or length of follow-up. We excluded case reports, case series, prospective and retrospective cohort studies, editorials, reviews and basic science papers.

Identification of trials

We used multiple strategies to identify potential eligible trials. MEDLINE, Embase and the Cochrane databases were systematically searched in Ovid up to and including Jan. 20, 2014. A health sciences librarian experienced in the conduct of systematic reviews assisted in developing and performing the search. We used medical subject headings, and Emtree headings and subheadings in various combinations, and supplemented with free text to increase sensitivity (Appendix 1, available at www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.140433/-/DC1). The search strategy was adapted in PubMed to search for articles published online ahead of print. Abstracts from recent major orthopedic and sports medicine conferences were reviewed. We consulted with experts in the field, manually reviewed the reference lists of articles that fulfilled the eligibility criteria and used the “related articles” feature in PubMed. Ongoing trials were identified from ClinicalTrials.gov.

Screening and assessment of eligibility

Two reviewers with methodologic and content expertise (M.K. and N.E.) independently screened all titles and abstracts for eligibility using a piloted electronic database (Microsoft Excel). All discrepancies were resolved by consensus. Duplicate articles were manually excluded. Both reviewers reviewed the full text of all studies identified by title and abstract screening to determine final eligibility.

Assessment of risk of bias

The same reviewers independently performed duplicate outcome-specific assessment of risk of bias using the Cochrane Collaboration’s tool for risk-of-bias assessment.¹⁹ When the issues bearing on the risk of bias were identical across outcomes, we made a single risk-of-bias assessment across outcomes.²¹

Extraction of data

Data were extracted independently and in duplicate by both reviewers using a piloted electronic data extraction form. If important data were unclear or not reported, attempts were made to contact the study authors for clarification.

Critical outcomes were determined to be patient-important outcomes related to pain, function and postintervention complications.^{22,23} Functional outcomes were measured by various disease-specific assessment scales.^{24–28} Postintervention pain was assessed using a visual analogue scale.²⁹

Statistical analysis

We calculated interobserver agreement for reviewer’s assessments of study eligibility with the Cohen κ coefficient.³⁰ Interobserver agreement for assessments of methodologic quality was calculated with the intraclass correlation coefficient. The κ and intraclass correlation coefficients were calculated using SPSS software (version 21.0, SPSS Inc.).

We used standardized mean differences (SMDs) to summarize outcome instruments that measured similar constructs.¹⁹ We pooled SMDs from individual trials to obtain the pooled estimate of effect for each outcome. The SMDs were weighted by sample size using the random effects model based on the inverse variance method.¹⁹ We transformed scores when required to ensure that higher scores indicated improved function in all cases.¹⁹ When standard deviations (SDs) were not available, they were calculated from alternative measures or were otherwise estimated from trials within the same comparison with similar scales, outcomes and periods.^{19,31} We extracted data from graphical representations when required. When knees were randomized in trials, they were treated as individual participants.

To improve interpretability, we converted SMD results to the Knee Injury and Osteoarthritis Outcome Score (KOOS).^{32,33} The KOOS evaluates patient outcomes over 5 domains. These include pain frequency and severity; symptoms; difficulty with activities of daily living; difficulty experienced with sport and recreational activities; and knee-related quality of life. This scoring system has been used extensively in this patient population, and has been validated and shown to be reliable and responsive to change.³⁴ The minimal important difference (MID) (i.e., the smallest difference that an informed patient would perceive as important enough to justify a change in management) is estimated to be 10 for the KOOS, which was converted to units of SD using the KOOS median SD.²⁷ The approximation of the SMD as a KOOS is carried out through the following formula: mean difference (KOOS units) = SMD \times (median SD of KOOS).³² A zone of clinical equiv-

alence based on the converted MID was projected onto the forest plots to aid interpretability.

Outcomes were dichotomized to short-term (< 6 mo) and long-term (< 2 yr) data. We pooled sham surgery and nonoperative treatment given the similar underlying conservative nature of the procedures. Complications were tabulated and presented descriptively.

To assess for publication bias, we constructed funnel plots that examined sample size versus exposure effect across included trials for functional outcome at 2-year follow-up.¹⁹ The forest and funnel plots were created with RevMan 5.2 (The Cochrane Collaboration).

Evaluation of heterogeneity and sensitivity analyses

We quantified heterogeneity using the χ^2 test for heterogeneity and the I^2 statistic.¹⁹ The I^2 statistic estimates the proportion of total variability between studies due to heterogeneity rather than chance alone. We considered I^2 less than 25% to indicate low heterogeneity and I^2 greater than 75% to indicate considerable heterogeneity.¹⁹ We developed a priori hypotheses to explore both potential artifactual and real differences of treatment effect across trials.³⁵ We planned for subgroup analysis based on year of study to account for potential evolution of surgical technique. Sensitivity analyses were planned for studies to investigate the effects of missing data and those trials at high risk of bias.³⁶

Results

Search results and study characteristics

The literature search identified 946 potentially relevant studies: 944 from the electronic search and 2 from the manual search. Seven RCTs ($n = 805$ patients) were eligible for inclusion in this review³⁷⁻⁴³ (Figure 1). No non-English articles were identified. The κ for overall agreement between reviewers for the final eligibility decision was 0.92 (95% confidence interval [CI] 0.85 to 1.00).

Of the 7 trials, 5 were conducted in Europe, 1 in the US³⁹ and 1 in South Korea.⁴¹ Four of the included trials were single-centre trials^{37,38,41,43} and 3 were multicentre trials^{39,40,42} (Table 1).

All eligible trials included patients with degenerative meniscal tears, documented by magnetic resonance imaging (MRI) or arthroscopy, in the setting of mild or no osteoarthritis. Sample sizes ranged from 8 to 330, and the total sample included 811 knees (805 patients). The mean age of patients was 56 (± 3.2) years. Age eligibility criteria from included studies ranged from 35 to 65 years when reported.^{37-40,42}

Two trials^{37,41} did not report crossovers to the surgical arm. Of the trials that did report cross-

over, Herrlin and colleagues³⁸ reported 27%, Katz and colleagues³⁹ reported 30%, Østerås and colleagues⁴² reported 0%, Vermesan and colleagues⁴³ reported 17%, and Sihvonen and colleagues⁴⁰ reported 7%. Six trials³⁷⁻⁴² documented patients who declined to participate, with rates ranging from 5%³⁹ to 40%.³⁸ Reasons for non-participation when described included preference for arthroscopic or conservative management, lack of symptoms while on surgical wait list and refusal due to time commitment.

Three trials specifically documented that no other surgical co-interventions were performed during the arthroscopic débridement procedure.³⁹⁻⁴¹ Five of the 7 included studies documented similar nonoperative rehabilitation programs between groups undergoing operative and nonoperative interventions.³⁷⁻⁴¹

Risk of bias

Only 1 included study was found to have a low risk of bias.⁴⁰ The remainder of the included trials were found to have uncertain to high risk of bias (Figure 2). Agreement between reviewers in the assessment of risk of bias was high (intra-class correlation coefficient 0.93, 95% CI 0.88 to 0.96). Appendix 2 (available at www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.140433/-/DC1)




Figure 1: Selection of randomized controlled trials (RCTs) for inclusion in the meta-analysis.

presents a summary of findings providing outcomes along with an evaluation of the quality of evidence based on the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach.²¹

Function

Arthroscopic débridement resulted in a significant improvement in short-term function across 6 trials^{37,39-43} involving a total of 805 patients (SMD 0.25, 95% CI 0.02 to 0.48) with moderate hetero-

Table 1: Studies included in the meta-analysis

Study	Year	Country	Patient characteristics		Treatment arm, no. of patients and description of treatment*		Major outcome measures	OA inclusion	Loss to follow-up
			Mean age, yr	Male sex, %	Conservative	Surgical			
Herrlin et al. ³⁷	2007	Sweden	55	61	43 Standardized exercise program for 8 wk	47 Arthroscopic meniscal débridement	KOOS, Lysholm Knee Scoring Scale, Tegner Activity Scale, VAS scores at 8 wk, 6 mo	Ahlbäck criteria grade 0-1	NR
Herrlin et al. ³⁸	2013	Sweden	55	60	49 Standardized exercise program for 8 wk	47 Arthroscopic meniscal débridement	Lysholm Knee Scoring Scale, Tegner Activity Scale, VAS scores (at rest and with activity) at 24, 60 mo	Ahlbäck criteria grade 0-1	C: 2/49 S: 2/47
Katz et al. ³⁹	2013	United States	58	43	169 Land-based, individualized physical therapy with progressive home exercise	161 Arthroscopic meniscal débridement	WOMAC-pf, KOOS pain scale, SF-36 physical activity scores at 3, 6, 12 mo	Kellgren-Lawrence grade 0-3	C: 2/169 S: 1/161
Østerås et al. ⁴²	2012	Norway	50	76	9 Exercise program 3 times/wk for 3 mo	8 Arthroscopic meniscal débridement	KOOS, VAS, HAD scores, quadriceps muscle strength at 3 mo	Kellgren-Lawrence grade 0-2	0
Sihvonen et al. ⁴⁰	2013	Finland	52	61	76 Sham surgical procedure	70 Arthroscopic meniscal débridement	Lysholm Knee Scoring Scale, WOMET, VAS, 15D, patient satisfaction scores at 2, 6, 12 mo	Kellgren-Lawrence grade 0-1	0
Yim et al. ⁴¹	2013	South Korea	56	21	52 NSAIDs, 3 wk supervised physical exercise followed by 8 wk home exercise program	50 Arthroscopic meniscal débridement	Lysholm Knee Scoring Scale, VAS, patient satisfaction, Tegner Activity Scale scores at 3, 6 mo, 1, 2 yr	Kellgren-Lawrence grade 0-1	C: 2/54 S: 4/54
Vermesan et al. ⁴³	2013	Romania	58	23	60 knees Interarticular steroid injection	60 knees Arthroscopic meniscal débridement	Oxford Knee Score at 1 mo, 1 yr	Kellgren-Lawrence grade 0-1	0

Note: 15D = a health-related quality-of-life scale made up of 15 dimensions on a scale of 0 (death) to 1 (full health), C = conservative treatment arm, HAD = Hospital Anxiety and Depression, KOOS = Knee Injury and Osteoarthritis Outcome Score, NR = not reported, NSAIDs = nonsteroidal anti-inflammatory drugs, OA = osteoarthritis, S = surgical treatment arm, SF-36 = Short Form 36, VAS = visual analogue scale, WOMAC-pf = Western Ontario and McMaster Universities Osteoarthritis Index — physical function, WOMET = Western Ontario Meniscal Evaluation Tool.
*Conservative to surgical crossover: Herrlin et al.,³⁷ NR; Herrlin et al.,³⁸ 27%; Katz et al.,³⁹ 30%; Østerås et al.,⁴² 0%; Sihvonen et al.,⁴⁰ 7%; Yim et al.,⁴¹ NR; Vermesan et al.,⁴³ 17%.

geneity ($p = 0.04$, $I^2 = 56\%$). This is equivalent to an estimated KOOS mean difference of 5.6 (95% CI 0.45 to 10.8). The KOOS was included from 3 trials,^{37,39,42} and scores from the Western Ontario Meniscal Evaluation Tool,⁴⁰ the Lysholm Knee Scoring Scale,⁴¹ and the Oxford Knee Score⁴³ were included from 1 trial each. This treatment effect failed to exceed the threshold of patient importance based on the MID (Figure 3).

We conducted an a priori subgroup analysis to evaluate trials reporting identical outcome measures and to evaluate trials by type of conservative treatment. Trials that used the KOOS (2 trials,^{37,42} $n = 107$ patients) did not show significant improvement in short-term function (SMD 0.03, 95% CI -0.35 to 0.41) and had low heterogeneity ($p = 0.86$, $I^2 = 0\%$). Removal of a study in which intra-articular steroid injection was given⁴³ in comparison with arthroscopic débridement decreased heterogeneity substantially (56% to 0%), which potentially accounts for the significant between-study variability with no effect on the pooled treatment effect (SMD 0.16, 95% CI 0.01 to 0.31).

Five trials^{38-41,43} involving a total of 794 patients or knees that evaluated long-term function following arthroscopic débridement did not show a significant improvement in function (SMD 0.07, 95% CI -0.10 to 0.23). Heterogeneity was low ($p = 0.28$, $I^2 = 20\%$). Scores from the Lysholm Knee Scoring Scale were pooled from 3 trials;^{38,40,41} KOOS³⁹ and Oxford Knee Score⁴³ values were pooled from the remaining 2 trials in this analysis (Figure 4). The result is equivalent to an estimated KOOS mean difference of 1.6 (95% CI -2.2 to 5.2), which failed to exceed the threshold of patient importance based on the MID.

Trials presenting scores from the Lysholm Knee Scoring Scale (3 trials,^{38,40,41} $n = 344$ patients) did not report significant results (SMD 0.00, 95% CI -0.22 to 0.21) and had low heterogeneity ($p = 0.46$, $I^2 = 0\%$).

Pain

Arthroscopic treatment did not improve short-term pain across 4 trials^{37,40-42} that reported short-term visual analogue scores from 355 patients (mean difference [MD] 0.20, 95% CI -0.67 to 0.26) with low heterogeneity ($p = 0.36$, $I^2 = 6\%$). Similarly, long-term pain after arthroscopic débridement across 3 trials^{38,40,41} involving 344 patients did not show a significant improvement in pain scores (MD -0.06, 95% CI -0.28 to 0.15) and had low heterogeneity ($p = 0.75$, $I^2 = 0\%$).

Adverse events

Two trials reported on adverse events: Sihvonen and colleagues⁴⁰ reported 1 infection in the group undergoing arthroscopic meniscal

débridement as compared with a sham procedure. Katz and colleagues³⁹ reported 3 serious adverse events in the group undergoing arthroscopic meniscal débridement and 2 in the group undergoing physical therapy.

Sensitivity analysis

We conducted a sensitivity analysis to investigate the effects of estimated missing SDs on long-term function and pain through the removal of studies that required estimated SDs. The results were not significant⁴¹ (SMD 0.10, 95% CI -0.09 to 0.29), and heterogeneity was low ($p = 0.23$, $I^2 = 31\%$). Sensitivity analysis related to sample size did not show a significant effect⁴² (SMD 0.26, 95% CI 0.02 to 0.51), and heterogeneity was substantial ($p = 0.03$, $I^2 = 64\%$).

Interpretation

Arthroscopic surgery for degenerative meniscal tears in the setting of mild or no concurrent osteoarthritis in middle-aged patients may have little, if any, effect on short-term (< 6 mo) and long-term (< 2 yr) outcomes in comparison with nonoperative management (Appendix 2).

The results of this meta-analysis are similar to those of recent trials by Moseley and colleagues⁸ and Kirkley and colleagues,⁹ which showed no

	Adequate sequence generation?	Allocation concealment?	Blinding?	Incomplete outcome data addressed?	Free of selective reporting?	Free of other bias?
Herrlin et al. ³⁷	?	?	-	?	?	+
Herrlin et al. ³⁸	+	+	-	+	-	+
Katz et al. ³⁹	+	+	-	+	+	+
Østerås et al. ⁴²	?	?	-	+	+	+
Sihvonen et al. ⁴⁰	+	+	+	+	+	+
Vermesan et al. ⁴³	?	?	-	?	?	?
Yim et al. ⁴¹	?	?	-	?	+	+

+ = low risk, ? = uncertain risk, - = high risk

Figure 2: Risk-of-bias assessment of randomized controlled trials included in the meta-analysis.

benefit of arthroscopic débridement or lavage for osteoarthritis. Our findings extend the evidence to show a lack of clinical efficacy in surgical débridement of meniscal tears in the setting of mild or no osteoarthritis.

The results of our investigation indicate that arthroscopic management did not exceed the MID in comparison with nonoperative management in both the short and long term. The MID is the smallest effect that an informed patient would perceive as valuable enough to justify a change in therapeutic management when weighing the anticipated benefits against the possible harms of an intervention.^{44,45} The MID concept has also been referred to as the minimal clinically important difference or the minimal clinically important improvement.⁴⁵ It can be estimated with an anchor-based approach (which correlates the score of interest with a known measure of clinical change) or a distribution-based approach (which suggests that one-half of an SD of a continuous outcome score constitutes a clinically

meaningful difference).²³ Although not without limitations, this tool aids clinicians in evaluating therapeutic options and determining whether significant outcomes will have clinically meaningful implications.²³ A limitation of this approach is that MIDs may be context-specific and may not be applicable across treatments or populations. Minimal important differences must therefore be defined for specific populations to provide useful guidance to users of these instruments.⁴⁵ The MID has not been clearly defined for the KOOS, and an estimated 10-point change in the scale is considered the MID based on half-SD methods, although research is ongoing.^{27,46-48} Future research is required to define a range of MID values for various clinical contexts to accurately identify true clinically important differences for all instruments measuring quality of life. Given these current limitations, the MID should be used as a supplementary tool for clinicians in determining the relevance of study findings.⁴⁵




Figure 3: Pooled short-term functional outcomes of conservative and surgical treatment. Red lines show a zone of clinical equivalence based on a minimal important difference of 10 on the Knee Injury and Osteoarthritis Outcome Score.^{37,39,40-43}
 Note: CI = confidence interval, SD = standard deviation, SMD = standardized mean difference.




Figure 4: Pooled long-term functional outcomes of conservative and surgical treatment. Red lines show a zone of clinical equivalence based on a minimal important difference of 10 on the Knee Injury and Osteoarthritis Outcome Score.^{38-41,43}
 Note: CI = confidence interval, SD = standard deviation, SMD = standardized mean difference.

Limitations

When data were unavailable despite attempts to contact the authors, we estimated SDs based on similar studies. A sensitivity analysis confirmed that this was unlikely to change the results of our study. Various outcome measures were combined in the pooled analysis; however, given their disease-specific similarities, we believe this was justified. Heterogeneity was low except for in the pooled SMD scores for functional outcomes at 6 months, which is potentially explained by variation in methods of conservative treatment. A funnel plot analysis suggested a low risk of publication bias (Appendix 3, available at www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.140433/-/DC1). A subgroup analysis related to year of publication was not required, because all included trials were conducted within the last 6 years.

Implications for practice and research

In the context of limited health care resources, clinicians must carefully select patients with degenerative meniscal pathology who would benefit from surgical intervention.^{2,49} Certain prognostic factors have been identified in the literature; for example, high levels of pain at baseline correlate with inferior patient-reported outcomes after arthroscopy.^{16,50} Assessment of a patient's quality of life is essential, given that abnormalities in anatomy and on MRI are not always correlated with symptoms.^{7,51} With limited evidence supporting arthroscopic meniscal débridement for degenerative meniscal tears in the setting of mild or no concomitant osteoarthritis, an initial trial of nonoperative interventions should play a large role for middle-aged patients.

Limited reporting outcomes and methods are highlighted in this review and have been identified as an important problem in the surgical literature.^{52,53} Five trials are currently registered on ClinicalTrials.gov, with 4 ongoing⁵⁴⁻⁵⁷ and 1 listed as completed⁵⁸ but unpublished. Results of these trials when available will further improve our confidence in the effect of treatment.

We identified a number of patients who declined to participate or who crossed over to a different treatment arm, which may have confounded results. Future research will be important to identify the prognosis of these patients. Additionally, economic evaluation is required to assess direct and indirect costs associated with arthroscopic meniscal débridement in comparison with various options for nonoperative treatment.⁵⁹

Future investigation into the impact of cartilage status, mechanical alignment, extent of meniscal damage, duration, severity and characteristics of symptoms, body mass index and baseline functional outcome scores may allow clinicians to fur-

ther determine who may benefit from arthroscopy in this population. Comparison of various rehabilitation protocols, adjunct modalities and injections will further define optimal initial nonoperative management. Studies into novel biological treatment options are in progress and may provide additional options for clinicians.⁶⁰

Conclusion

This systematic review and meta-analysis showed moderate evidence to suggest that there is no benefit to arthroscopic meniscal débridement for degenerative meniscal tears in comparison with nonoperative or sham treatment options for middle-aged patients with mild or no concomitant osteoarthritis. Future research is required to identify how indications and patient selection influence outcomes following surgical and conservative treatment.

References

1. *Arthroscopy. AOSSM sports tips*. 2008. American Orthopaedic Society for Sports Medicine. Available: www.sportsmed.org/uploadedFiles/Content/Patient/Sports_Tips/ST%20Arthroscopy%2008.pdf (accessed 2014 Feb. 16).
2. Kim S, Bosque J, Meehan JP, et al. Increase in outpatient knee arthroscopy in the United States: a comparison of National Surveys of Ambulatory Surgery, 1996 and 2006. *J Bone Joint Surg Am* 2011;93:994-1000.
3. Cullen KA, Hall MJ, Golosinskiy A. Ambulatory surgery in the United States, 2006. *Natl Health Stat Report* 2009;(11):1-25.
4. Hawker G, Guan J, Judge A, et al. Knee arthroscopy in England and Ontario: patterns of use, changes over time, and relationship to total knee replacement. *J Bone Joint Surg Am* 2008;90:2337-45.
5. Health Quality Ontario. Arthroscopic lavage and debridement for osteoarthritis of the knee: an evidence-based analysis. *Ont Health Technol Assess Ser* 2005;5:1-37.
6. Buchbinder R. Meniscectomy in patients with knee osteoarthritis and a meniscal tear? *N Engl J Med* 2013;368:1740-1.
7. Marx RG. Arthroscopic surgery for osteoarthritis of the knee? *N Engl J Med* 2008;359:1169-70.
8. Moseley JB, O'Malley K, Petersen NJ, et al. A controlled trial of arthroscopic surgery for osteoarthritis of the knee. *N Engl J Med* 2002;347:81-8.
9. Kirkley A, Birmingham TB, Litchfield RB, et al. A randomized trial of arthroscopic surgery for osteoarthritis of the knee. *N Engl J Med* 2008;359:1097-107.
10. Mäntyselkä P, Kumpusalo E, Ahonen R, et al. Pain as a reason to visit the doctor: a study in Finnish primary health care. *Pain* 2001; 89:175-80.
11. Bonamo JJ, Kessler KJ, Noah J. Arthroscopic meniscectomy in patients over the age of 40. *Am J Sports Med* 1992;20:422-8, discussion 428-9.
12. Baumgaertner MR, Cannon WD Jr, Vittori JM, et al. Arthroscopic débridement of the arthritic knee. *Clin Orthop Relat Res* 1990;(253):197-202.
13. Bert JM, Maschka K. The arthroscopic treatment of unicompartmental gonarthrosis: a five-year follow-up study of abrasion arthroplasty plus arthroscopic débridement and arthroscopic débridement alone. *Arthroscopy* 1989;5:25-32.
14. Gross DE, Brenner SL, Esformes I, et al. Arthroscopic treatment of degenerative joint disease of the knee. *Orthopedics* 1991;14: 1317-21.
15. Börjesson M, Robertson E, Weidenhielm L, et al. Physiotherapy in knee osteoarthritis: effect on pain and walking. *Physiother Res Int* 1996;1:89-97.
16. Rathleff CR, Cavallius C, Jensen HP, et al. Successful conservative treatment of patients with MRI-verified meniscal lesions. *Knee Surg Sports Traumatol Arthrosc* 2013 Apr. 11. [Epub ahead of print].
17. Jevsevar DS, Brown GA, Jones DL, et al. The American Academy of Orthopaedic Surgeons evidence-based guideline on: treatment of osteoarthritis of the knee, 2nd edition. *J Bone Joint Surg Am* 2013;95:1885-6.
18. Kieser C. A review of the complications of arthroscopic knee surgery. *Arthroscopy* 1992;8:79-83.

19. Higgins JPT, Green S, editors. Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011.
20. Moher D, Liberati A, Tetzlaff J, et al. PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009;62:1006-12.
21. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64:383-94.
22. Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines: 2. Framing the question and deciding on important outcomes. *J Clin Epidemiol* 2011;64:395-400.
23. Zlowodzki M, Bhandari M. Outcome measures and implications for sample-size calculations. *J Bone Joint Surg Am* 2009; 91(Suppl 3):35-40.
24. Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res* 1985;(198):43-9.
25. Lysholm J, Gillquist J. Evaluation of knee ligament surgery results with special emphasis on use of a scoring scale. *Am J Sports Med* 1982;10:150-4.
26. Dawson J, Fitzpatrick R, Murray D, et al. Questionnaire on the perceptions of patients about total knee replacement. *J Bone Joint Surg Br* 1998;80:63-9.
27. Roos EM, Lohmander LS. Knee injury and Osteoarthritis Outcome Score (KOOS): from joint injury to osteoarthritis. *Health Qual Life Outcomes* 2003;1:64.
28. Kirkley A, Griffin S, Whelan D. The development and validation of a quality of life-measurement tool for patients with meniscal pathology: the Western Ontario Meniscal Evaluation Tool (WOMET). *Clin J Sport Med* 2007;17:349-56.
29. Bhandari M, Petrisor B, Schemitsch E. Outcome measurements in orthopedic. *Indian J Orthop* 2007;41:32-6.
30. Fleiss JL. *Statistical methods for rates and proportions*. 2nd ed. New York: John Wiley and Sons; 1981.
31. Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Med Res Methodol* 2005;5:13.
32. Thorlund K, Walter SD, Johnston BC, et al. Pooling health-related quality of life outcomes in meta-analysis — a tutorial and review of methods for enhancing interpretability. *Res Synth Methods* 2011;2:188-203.
33. Guyatt GH, Thorlund K, Oxman AD, et al. GRADE guidelines: 13. Preparing summary of findings tables and evidence profiles-continuous outcomes. *J Clin Epidemiol* 2013;66:173-83.
34. Collins NJ, Misra D, Felson DT, et al. Measures of knee function: International Knee Documentation Committee (IKDC) Subjective Knee Evaluation Form, Knee Injury and Osteoarthritis Outcome Score (KOOS), Knee Injury and Osteoarthritis Outcome Score Physical Function Short Form (KOOS-PS), Knee Outcome Survey Activities of Daily Living Scale (KOS-ADL), Lysholm Knee Scoring Scale, Oxford Knee Score (OKS), Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), Activity Rating Scale (ARS), and Tegner Activity Score (TAS). *Arthritis Care Res (Hoboken)* 2011;63(Suppl 11):S208-28.
35. Glasziou PP, Sanders SL. Investigating causes of heterogeneity in systematic reviews. *Stat Med* 2002;21:1503-11.
36. Higgins JP, White IR, Wood AM. Imputation methods for missing outcome data in meta-analysis of clinical trials. *Clin Trials* 2008;5:225-39.
37. Herrlin S, Hallander M, Wange P, et al. Arthroscopic or conservative treatment of degenerative medial meniscal tears: a prospective randomised trial. *Knee Surg Sports Traumatol Arthrosc* 2007;15:393-401.
38. Herrlin SV, Wange PO, Lapidus G, et al. Is arthroscopic surgery beneficial in treating non-traumatic, degenerative medial meniscal tears? A five year follow-up. *Knee Surg Sports Traumatol Arthrosc* 2013;21:358-64.
39. Katz JN, Brophy RH, Chaisson CE, et al. Surgery versus physical therapy for a meniscal tear and osteoarthritis. *N Engl J Med* 2013; 368:1675-84.
40. Sihvonen R, Paavola M, Malmivaara A, et al. Finnish Degenerative Meniscal Lesion Study (FIDELITY) Group. Arthroscopic partial meniscectomy versus sham surgery for a degenerative meniscal tear. *N Engl J Med* 2013;369:2515-24.
41. Yim JH, Seon JK, Song EK, et al. A comparative study of meniscectomy and nonoperative treatment for degenerative horizontal tears of the medial meniscus. *Am J Sports Med* 2013;41:1565-70.
42. Østerås H, Østerås B, Torstensen TA. Medical exercise therapy, and not arthroscopic surgery, resulted in decreased depression and anxiety in patients with degenerative meniscus injury. *J Bodyw Mov Ther* 2012;16:456-63.
43. Vermesan D, Prejbeanu R, Laitin S, et al. Arthroscopic debridement compared to intra-articular steroids in treating degenerative medial meniscal tears. *Eur Rev Med Pharmacol Sci* 2013;17:3192-6.
44. Schünemann HJ, Guyatt GH. Commentary — goodbye M(C)ID! Hello MID, where do you come from? *Health Serv Res* 2005; 40:593-7.
45. Bannuru RR, Vaysbrot EE, McIntyre LF. Did the American Academy of Orthopaedic Surgeons osteoarthritis guidelines miss the mark? *Arthroscopy* 2014;30:86-9.
46. Singh JA, Luo R, Landon GC, et al. Reliability and clinically important improvement thresholds for osteoarthritis pain and function scales: a multicenter study. *J Rheumatol* 2014;41:509-15.
47. Norman GR, Sloan JA, Wyrwich KW. Interpretation of changes in health-related quality of life: the remarkable universality of half a standard deviation. *Med Care* 2003;41:582-92.
48. Engelhart L, Nelson L, Lewis S, et al. Validation of the Knee Injury and Osteoarthritis Outcome Score subscales for patients with articular cartilage lesions of the knee. *Am J Sports Med* 2012;40:2264-72.
49. Tarride JE, Haq M, O'Reilly DJ, et al. The excess burden of osteoarthritis in the province of Ontario, Canada. *Arthritis Rheum* 2012;64:1153-61.
50. Mallen CD, Peat G, Thomas E, et al. Prognostic factors for musculoskeletal pain in primary care: a systematic review. *Br J Gen Pract* 2007;57:655-61.
51. Hack K, Di Primio G, Rakhra K, et al. Prevalence of cam-type femoroacetabular impingement morphology in asymptomatic volunteers. *J Bone Joint Surg Am* 2010;92:2436-44.
52. Poolman RW, Struijs PA, Krips R, et al. Does a "Level I Evidence" rating imply high quality of reporting in orthopaedic randomised controlled trials? *BMC Med Res Methodol* 2006;6:44.
53. Jacquier I, Boutron I, Moher D, et al. The reporting of randomized clinical trials using a surgical intervention is in need of immediate improvement: a systematic review. *Ann Surg* 2006;244:677-83.
54. Department of Veterans Affairs. Prospective trial of arthroscopic meniscectomy for degenerative meniscus tears. In: ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). 2000-[cited 2014 Feb. 15]. Available: <http://clinicaltrials.gov/ct2/show/NCT01931735> (accessed 2014 Feb. 15). NLM identifier: NCT01931735.
55. Oslo University Hospital. Surgical or exercise therapy on patients with degenerative meniscus tears. In: ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). 2000-[cited 2014 Feb. 15]. Available: <http://clinicaltrials.gov/ct2/show/NCT01002794> (accessed 2014 Feb. 15). NLM identifier: NCT01002794.
56. Slagelse Hospital. The benefit of arthroscopic partial meniscectomy in middle-aged patients (SLAMSHAM). In: ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). 2000. Available: <http://clinicaltrials.gov/ct2/show/study/NCT01264991> (accessed 2014 Feb. 15). NLM identifier: NCT01264991.
57. Onze Lieve Vrouwe Gasthuis. Early Surgery Versus Conservative Therapy for Meniscal Injuries in Older Patients (ESCAPE). In: ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). 2000. Available: <http://clinicaltrials.gov/ct2/show/study/NCT01850719> (accessed 2014 Feb. 15). NLM identifier: NCT01850719.
58. University Hospital, Linköeping. Is a knee arthroscopy of any benefit for the middleaged patient with meniscal symptoms? In: ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). 2000. Available: <http://clinicaltrials.gov/ct2/show/study/NCT01288768> (accessed 2014 Feb. 15). NLM identifier: NCT01288768.
59. van de Graaf VA, de Gast A, Poolman RW. [Arthroscopic meniscectomy: Does it make sense in patients older than 45?]. [Article in Dutch] *Ned Tijdschr Geneesk* 2013;157:A6865.
60. Kon E, Filardo G, Drobnic M, et al. Non-surgical management of early knee osteoarthritis. *Knee Surg Sports Traumatol Arthrosc* 2012;20:436-49.

Affiliations: Division of Orthopaedic Surgery (Khan, Evaniew, Ayeni, Bhandari), Department of Surgery, McMaster University, Hamilton, Ont.; MedSport, Department of Orthopaedic Surgery (Bedi), University of Michigan, Ann Arbor, Mich.; Department of Clinical Epidemiology and Biostatistics (Bhandari), McMaster University, Hamilton, Ont.

Contributors: Moin Khan and Nathan Evaniew contributed to the review and quality assessment of all included studies and conducted the statistical analysis. Asheesh Bedi, Olufemi Ayeni and Mohit Bhandari reviewed the data. Mohit Bhandari contributed to the methods and the statistical analysis. Moin Khan and Nathan Evaniew drafted the article, which Asheesh Bedi, Olufemi Ayeni and Mohit Bhandari revised. All of the authors approved the version submitted for publication and agreed to act as guarantors of the work.