

Polypharmacy: prevention and management

We have followed the *CMAJ* series of case-based practice articles on polypharmacy with great interest and we commend the authors^{1,2} for drawing attention to this issue. As the population of older Canadians with multiple chronic conditions grows, the challenge of managing medications in these patients becomes more pressing.

Farrell and colleagues^{1,2} have presented two instructive cases of senior patients with multiple chronic conditions, complex medication regimens and multiple prescribers. We wish to emphasize the essential role of the interprofessional team in making sense of the complexities inherent to such cases. In both the cases^{1,2} the patients benefited greatly from referral to a local geriatric day hospital. Unfortunately, current demand for such specialized services far outstrips supply. Consequently, much of the management of complex medication regimens is performed in primary care. This is a time-consuming process, as Frank³ has indicated, and is not well suited to the usual 10- to 15-minute family-physician visit.

Necessity being the mother of invention, we developed a new primary care model designed specifically for older patients with complex health and medication needs. The IMPACT clinic⁴ features an extended 90-minute visit, during which an interprofessional team conducts a comprehensive 360-degree assessment and co-creates, with the patient and family, a plan of care that is mutually agreed upon. Review of medication appropriateness and reconciliation is a vitally important component of the clinic, and all IMPACT patients and their families are provided with an up-to-date, user-friendly medication list.

Given the high frequency of change in complex regimens, which are often driven by visits to multiple prescribers, we believe that primary care is the appropriate setting for ongoing medication management in complex patients. The series by Farrell and colleagues^{1,2}

also underscores the importance of applied research to develop, implement and evaluate management tools for complex medication regimens.

C. Shawn Tracy MSc, Jonathan Fuller BMSc, Ross E.G. Upshur MD MSc
Research scientist (Tracy), Medical Director (Upshur), Bridgepoint Hospital, and University of Toronto (Tracy, Fuller, Upshur), Toronto, Ont.

References

- Farrell B, Merkley VF, Thompson W. Managing polypharmacy in a 77-year-old woman with multiple prescribers. *CMAJ* 2013;185:1240-5.
- Farrell B, Monahan A, Thompson W. Revisiting medication use in a frail 93-year-old man experiencing possible adverse effects. *CMAJ* 2014;186:445-9.
- Frank C. Deprescribing: a new word to guide medication review. *CMAJ* 2014;186:407-8.
- Tracy CS, Bell SH, Nickell LA, et al. The IMPACT clinic: an innovative model of interprofessional primary care for elderly patients with complex health-care needs. *Can Fam Physician* 2013;59:e148-55.

CMAJ 2014. DOI:10.1503/cmaj.114-0082

The authors respond

We thank Tracy and colleagues for their supportive letter¹ in response to our series of case-based practice articles on polypharmacy.^{2,3} Their letter accentuates the challenges of complex, ever-changing medication regimens, numerous comorbidities, multiple prescribers and time constraints in the effort to effect positive medication change. We applaud Tracy and colleagues for their initiative.

We agree that more tools are needed to support primary care physicians in their work with seniors. Not all physicians have access to a specialized interdisciplinary team, nor are they able to afford a significant block of time to address all the issues at hand. We hope that our case series (a total of eight case reports and two commentaries in *Canadian Family Physician* and the *Canadian Pharmacists Journal*)⁴⁻¹¹ will help physicians to prioritize medication changes and to realize that these changes do not need to happen all at once. In emphasizing the supporting roles of the various health professionals on our team, we hope to assist primary care physicians to determine whether some functions could be carried out by existing team members. Physicians should also be aware of local community supports, such as physiother-

apy clinics with fall-prevention programs, when access to a specialized geriatric team is not available. We recognize that this is a work in progress.

Salima Shamji MD, Barbara Farrell BScPhm PharmD, Ann Monahan MD, Véronique French Merkley MD
Bruyère Continuing Care, Ottawa, Ont.

References

- Tracy SC, Fuller J, Upshur REG. On prevention and management of polypharmacy [letter]. *CMAJ* 2014;183:1321.
- Farrell B, Merkley VF, Thompson W. Managing polypharmacy in a 77-year-old woman with multiple prescribers. *CMAJ* 2013;185:1240-5.
- Farrell B, Monahan A, Thompson W. Revisiting medication use in a frail 93-year-old man experiencing possible adverse effects. *CMAJ* 2014;186:445-9.
- Farrell B, Shamji S, Monahan A, et al. Reducing polypharmacy in the elderly: Cases to help you rock the boat. *Can Pharm J (Ott)* 2013;146:243-4.
- Farrell B, Shamji S, Monahan A, et al. Clinical vignettes to help you prescribe medications in elderly patients: introduction to the polypharmacy case series. *Can Fam Physician* 2013;59:1257-8.
- Farrell B, Shamji S, Dalton D. Managing chronic disease in the frail elderly: more than just adhering to clinical guidelines. *Can Pharm J (Ott)* 2014;147:89-96.
- Farrell B, Eisener-Parsche P, Dalton D. Turning over the rocks: the role of anticholinergics and benzodiazepines in cognitive decline and falls. *Can Fam Physician* 2014;60:345-50.
- Farrell B, Monahan A, Ingar N. Identifying and managing drug-related causes of common geriatric symptoms. *Can Fam Physician* 2014;60:147-53.
- Farrell B, French Merkley V, Ingar N. Reducing pill burden and helping with medication awareness to improve adherence. *Can Pharm J (Ott)* 2013;146:262-9.
- Farrell B, Shamji S, Ingar N. Reducing fall risk while managing pain and insomnia: addressing polypharmacy in an 81-year-old woman. *Can Pharm J (Ott)* 2013;146:335-41.
- Farrell B, Shamji S, Ingar N. Reducing fall risk while managing hypotension, pain and poor sleep in an 83-year-old woman. *Can Fam Physician* 2013;59:1300-5.

CMAJ 2014. DOI:10.1503/cmaj.114-0083

Dental profession fails to meet needs of disabled Canadians

Kelsall and O'Keefe¹ emphasize the poor health implications for seniors caused by their inability to pay for necessary dental treatment, but disabled Canadians are also seriously affected by inequitable access to dental care.

Provincial-government dental plans for the disabled have stagnated for years even as dental fees have increased annually. Some government plans now pay only 50%–60% of typi-