

## CLINICAL IMAGES

## Bilateral central scotomas following laser pointer misuse

Mark Xu HBSc, Tony Lin MD




Figure 1: Colour fundus photographs of the right (A) and left (B) eye of a 12-year-old boy with bilateral central blind spots. \*Pigmentary changes in the macula and fovea of both eyes.

A 12-year-old white boy presented with a 3-day history of bilateral central blind spots after staring directly into the beam of a red laser pointer for about 1 minute with each eye. His Snellen visual acuity was 20/25 in his right eye and 20/20 in his left eye. Fundoscopy showed foveal and parafoveal pigmentary changes in both maculae (Figure 1). Automated visual field testing of the central 10 degrees was unremarkable, which suggested that the retinal damage affected less than 2 degrees of the visual angle. Optical coherence tomography of the maculae showed bilateral damage to the photoreceptors and retinal pigment epithelium in the fovea (Appendix 1, available at [www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130975/-/DC1](http://www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130975/-/DC1)). Treatment options were limited, and we decided not to actively intervene. Re-evaluation at 6 weeks showed bilateral changes to the retinal pigment epithelium, consistent with scarring of the foveal burns. The patient's vision returned to normal at 4 months, with no improvement in retinal findings.

Laser pointers have been implicated in eye injuries resulting in permanent vision loss, particularly among children.<sup>1</sup> It is prohibited to sell, lease or import hazardous, high-powered laser pointers (> 5 mW) in Canada.<sup>2</sup> Our patient had been able to obtain a laser pointer with a

measured output of 48 mW as a gift purchased overseas. His retinal injury likely resulted from a combination of a high-powered laser and long duration of exposure.

Proximity of injury to the fovea is the most important determinant of functional visual impairment.<sup>3</sup> Foveal and parafoveal burns directly reduce visual acuity, whereas burns more distant from the fovea may produce a scotoma.<sup>3</sup> Vision generally improves over days to months, and prognosis is excellent if the fovea is uninvolved.<sup>4</sup> Visual cortex remapping may allow perceptual fill-in and improvement of the scotomas, as seen in computational modelling.<sup>5</sup>

## References

1. FDA safety notification: risk of eye and skin injuries from high-powered, hand-held lasers used for pointing or entertainment. Silver Spring (MD): US Food and Drug Administration; 2010. Available: [www.fda.gov/MedicalDevices/Safety/AlertsandNotices/ucm237129.htm](http://www.fda.gov/MedicalDevices/Safety/AlertsandNotices/ucm237129.htm) (accessed 2013 Oct. 11)
2. Hand-held lasers or laser pointers may cause permanent vision damage and burns. Ottawa (ON): Health Canada; 2012. Available: [www.healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2012/15008a-eng.php](http://www.healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2012/15008a-eng.php) (accessed 2013 Oct. 11)
3. Barkana Y, Belkin M. Laser eye injuries. *Surv Ophthalmol* 2000; 44:459-78.
4. Mainster MA, Stuck BE, Brown J Jr. Assessment of alleged retinal laser injuries. *Arch Ophthalmol* 2004;122:1210-7.
5. McManus JN, Ullman S, Gilbert CD. A computational model of perceptual fill-in following retinal degeneration. *J Neurophysiol* 2008;99:2086-100.

**Affiliations:** Schulich School of Medicine & Dentistry, Western University, London, Ont.

**Competing interests:** None declared.

This article has been peer reviewed.

The authors have obtained patient consent.

## Acknowledgement:

The authors thank Alain A. Proulx for his expertise on the case.

## Correspondence to:

Mark Xu,  
mxu2014@meds.uwo.ca

*CMAJ* 2014, DOI:10.1503/cmaj.130975