| v ul lubic | icb y diagnostic couc(s) | ICD IO diagnostic couc(s) | i nysicians service | 1 locedure codes |
|--|------------------------------|-----------------------------|---------------------|---|
| | | | claims (OHIP) codes | |
| Exclusion criteria: pre- | 570.x-573.x, 456.0-456.21, | K70.X-77.X, I85.X, I86.4, | Diagnostic codes: | CCP: 62.x |
| existing liver disease or an | 996.82, 155.0, 155.2, 230.8, | I98.2, R17.X, B15,X-19.X, | 571, 573, 155, 230, | CCI: 10A13LAAG, 10A13LAW2, 10A13LAW3, |
| $alcohol$ -related $condition^{\dagger}$ | 782.4, 070.x, 291.x, 303.x, | Z72.1, Z13.3, F10.x, G62.1, | 787, 573, 070, 303, | 10A13LAX7, 10A13LAXXE, 10A13LAXXN, |
| | 305.0-305.03, 357.5, 425.5, | I42.6, K29.2, P04.3, R78.X, | 291 | 10A16HAT9, 10A35H.X, 10A52H.X, 10A52LA, |
| | 535.3, 535.31, 760.71, | T51.X, Z71.4,Z50.2, G31.2, | | 10A58WLXX.X, 10A58LAXXK, 10A59DAAD, |
| | 790.3, 977.3, 980.x V42.7, | Z86.4 | Fee codes: E765, | 10A59DAAG, 10A59DAGX, 10A59DAX7, |
| | V1007, V113, V79.1 | | G254, S275, S294, | 10A59HAX7, 10A59LAAD, 10A59LAAG, |
| | | | S295, Z551, Z554 | 10A59LAGX, 10A59LAX7, 10A74LA, |
| | | | | 10A85LAXXK, 10A85VCXXK, 10A85WLXX.X, |
| | | | | 10A87LA.X, 20A71.X, |
| Censoring criteria: viral | 070.x, 573.1 | B15.X-19.X | 070 | |
| hepatitis, alcoholic liver | 571.0-571.3 | K70, K71.1 | | |
| disease, biliary cirrhosis or | 571.6 | K74.3-K74.5 | | |
| toxic liver failure | | | | |
| Outcome: serious liver disease | | | | |
| Liver cirrhosis | 571.15 | K71.7, K74.0-K74.2, K74.6 | | |
| Liver failure and its | 570.0, 572.2-572.8, 456.0- | K72, K76.6, K76.7 | | |
| sequelae | 456.21 | | | |
| Liver transplant | V42.7, 996.82 | | Fee codes: | CCP: 62.4x |

ICD-9 diagnostic code(s) ICD-10 diagnostic code(s) Physicians' service

Procedure codes*

CCI: 10A85LAXXK, 10A85VCXXK, 10A85WLXXJ, 10A85WLXXJ

Appendix l (as supplied by the author). Diagnostic and procedure codes used to identify study variables

*Indicates coding by Canadian Classification of Procedures (corresponding to ICD-9 years) or Canadian Classification of Interventions (corresponding to ICD-10 years). **Based on the algorithm by Tu et al: 2 physician billing claims or 1 hospital discharge with a diagnosis of hypertension in a 2-year period¹. CCP Canadian Classification of Procedures; CCI Canadian Classification of Health Interventions; OHIP Ontario Health Insurance Plan

G254, S294, S295

--

--

--

278

272

401-403

[†] Includes alcohol abuse and medical conditions related to alcohol (e.g. alcoholic liver disease, gastritis, cardiomyopathy, dementia and polyneuropathy)

I10.X, I11.X, I12.X, I13.X,

Reference

Obesity

Dyslipidemia

Metabolic syndrome features

Arterial hypertension**

278.0

272.0-272.5

401.X-405.X

Variable

1. Tu K, Campbell NR, Chen ZL, et al. Accuracy of administrative databases in identifying patients with hypertension. Open Med 2007;1:e18-26.

E66.X

E78.X

I15.X