

Appendix 2 (as supplied by the authors): Table describing to whom clinical actions were directed (based on 149 reports)

<u>Health care provider, total times engaged (n)</u>	<u>N of participants (%)</u>
Most responsible physician/resident, n=132	
No involvement (n, %)	64 (43.0%)
At least once (n, %)	85 (57.0%)
Occupational therapist, n=104	
No involvement (n, %)	96 (64.4%)
At least once (n, %)	53 (35.6%)
Dietitian, n=71	
No involvement (n, %)	103 (69.1%)
At least once (n, %)	46 (30.9%)
Other, n=52	
No involvement	114 (76.5%)
At least once	35 (23.5%)
Physiotherapist, n=47	
No involvement (n, %)	126 (84.6%)
At least once (n, %)	23 (15.4%)
Pharmacist, n=35	
No involvement (n, %)	122 (81.9%)
At least once (n, %)	27 (18.1%)
In-clinic group, n=25	
No involvement (n, %)	128 (85.9%)
At least once (n, %)	21 (14.1%)
System navigator, n=19	
No involvement	136 (91.3%)
At least once	13 (8.7%)
Nurse practitioner, n=17	
No involvement	138 (92.6%)
At least once	11 (7.4%)
Co-booked, n=9	
No involvement	141 (94.6%)
At least once	8 (5.4%)
Community program, n=7	
No involvement (n, %)	143 (96.0%)
At least once (n, %)	6 (4.0%)
Social worker, n=7	
No involvement (n, %)	145 (97.3%)
At least once (n, %)	4 (2.7%)
External specialist, n=5	
No involvement (n, %)	144 (96.6%)
At least once (n, %)	5 (3.4%)
Geriatrician/ Geriatrics huddle, n=2	
No involvement (n, %)	147 (98.7%)
At least once (n, %)	2 (1.3%)
Psychologist, n=0	
No involvement (n, %)	149 (100%)