

Paula M. Brauer PhD

Family Relations and Applied Nutrition,
University of Guelph, Guelph, Ont.

References

1. de Gara C, Kanji A, Karmali S, et al. Canadian Task Force obesity guidelines are unbalanced [letter]. *CMAJ* 2015;187:827.
2. Brauer P, Connor Gorber S, Shaw E, et al. Recommendations for prevention of weight gain and use of behavioural and pharmacologic interventions to manage overweight and obesity in adults in primary care. *CMAJ* 2015;187:184-95.
3. Peirson L, Douketis J, Ciliska D, et al. Treatment for overweight and obesity in adult populations: a systematic review and meta-analysis. *CMAJ Open* 2015;2:E306-17.

CMAJ 2015. DOI:10.1503/cmaj.1150051

Let's talk chronic heart failure

We applaud *CMAJ's* efforts at medical education via their "Five things to know about ..." series. However, we believe that a recent paper by Moayedi and Kobulnik¹ falls somewhat short.

Specifically, heart failure is predominantly a disease of the elderly.² Elderly patients are often not candidates for mechanical circulatory support or cardiac transplantation. Although there are options such as cardiac resynchronization devices and implantable cardioverter defibrillators, the benefits of these therapies may be attenuated in the elderly.³ Therefore, we are generally left with medical management options for this patient population.

Although the authors point out that there are some promising new medications on the horizon, at present, patients with heart-failure face five-year mortality rates of about 50%.² This prognosis is worse than that of many patients with cancer.⁴ Despite this, physicians rarely encourage advance-care planning or discuss goals of care with their patients with heart failure.⁵ The Choosing Wisely Canada initiative recommends not delaying these conversations.⁶ We need to start having these discussions with our patients with heart failure. What better way to do this than to

include it in your "five-things-to-know-about-heart-failure" list?

Michael Slawnych MD PhD, Nakul Sharma MD, Debra Isaac MD, Jessica Simon MB ChB

Division of Cardiology, Libin Cardiovascular Institute (Slawnych, Sharma, Isaac); Division of Palliative Medicine (Simon), Department of Oncology, University of Calgary, Calgary, Alta.

References

1. Moayedi Y, Kobulnik J. Chronic heart failure with reduced ejection fraction. *CMAJ* 2015;187:518.
2. Go AS, Mozaffarian D, Roger VL, et al. Heart disease and stroke statistics — 2013 update: a report from the American Heart Association. *Circulation* 2013;127:e6-e245.
3. Hess PL, Al-Khatib SM, Han JY, et al. Survival benefit of the primary prevention implantable cardioverter-defibrillator among older patients. Does age matter? An analysis of pooled data from 5 clinical trials. *Circ Cardiovasc Qual Outcomes*. 2015;8:179-86.
4. Adler ED, Goldfinger JZ, Kalman J, et al. Palliative care in the treatment of advanced heart failure. *Circulation* 2009;120:2597-606.
5. Doctors reluctant to discuss end-of-life care with heart failure patients. American Heart Association meeting report, abstract 352. Dallas: American Heart Association; 2014. Available: <http://newsroom.heart.org/news/doctors-reluctant-to-discuss-end-of-life-care-with-heart-failure-patients?preview=5988>.
6. Palliative care: five things physicians and patients should question. Toronto: Choosing Wisely Canada; 2015. Available: www.choosingwiselycanada.org/recommendations/palliative-care/.

CMAJ 2015. DOI:10.1503/cmaj.1150052

Time to rethink EMRs

Although I agree with most of Hall's points,¹ I fear he has drawn a false analogy. The United States Health Information Technology for Economic and Clinical Health (HITECH) Act of 2009 was enacted to mobilize information technology to reduce skyrocketing costs associated with the health care system.²

Hall¹ states that the distracting effect of electronic medical records (EMRs) on the physician-patient interaction is well known. This is usually because of the usability of the EMR interface. Most systems are not designed for the patient encounter, and vendors are reluctant to customize their systems for physicians' workflow. It is also well

known that EMRs neither reduce health care costs nor increase efficiency. Poor usability decreases efficiency and frustrates the user.

The actual benefits of EMRs are likely unmeasurable. Recently, the American Medical Informatics Association has called for a re-evaluation of economic analyses in health information technology,³ with a focus on quality and patient-safety benefits. Electronic medical records can improve patient care even without cost reductions. I believe it would be a patient care disaster if we lost our provincial EMRs to save money.

Finally, I echo Hall's comments about lobbying administrators and politicians about the use of EMRs. We are past the tipping point but, as an important user group, it is vital that we continue to loudly advocate for usable systems that will allow increased efficiencies and not diminish the patient-physician relationship.

Darren A. Hudson MSc MD

School of Health Information Science,
University of Victoria, Victoria, BC

References

1. Hall T. Time to rethink EMRs. *CMAJ* 2015;187:602.
2. Crownover K. Implications of the HITECH Act of 2009. *Caring* 2009;28:40-2.
3. Riskin L, Koppel R, Riskin D. Re-examining health IT policy: What will it take to derive value from our investment? *J Am Med Inform Assoc* 2015;22:459-64.

Competing interests: Darren Hudson is the associate medical director for the eCritical Alberta Critical Care Information System, and he has a masters degree in health information science from the University of Victoria, Victoria, BC, where he holds an academic post.

CMAJ 2015. DOI:10.1503/cmaj.1150053

Letters to the editor

In submitting a letter, you automatically consent to having it appear online/in print. All letters accepted for print will be edited for space and style. See www.cmaj.ca for full versions and competing interests.