



## Show me the proof, Dr. Avery

The fall 1998 meeting during which US editorial writers met Dr. Granger Avery and Michael Decter reflects the abyss separating those who want more private care and those who support medicare's principles and see solutions not in its dissolution but in better management.<sup>1</sup>

Avery, a well-known devotee of expanded privatization, repeated statements that have become part of the mantra of privatizers. When these are used by those who favour a parallel private system, they should either be defended with objective data or discounted.<sup>2-4</sup>

For instance, Avery should present the data to support his statement that Canadians spend \$1 billion annually buying health care in the US, not because they fall ill while visiting but because they have specifically sought care that is unavailable in Canada or lack trust in the Canadian system. Given that this number is quoted so often, it should not be too difficult for Avery to present the source of his information.

His contention that Canada, Cuba and North Korea are the only countries with single-payer systems is completely misleading. The Scandinavian countries and The Netherlands may have various sources of payment for health care services, but — irrespective of the source of payment — access to the system is the same, other than for marginal services such as hospital accommodation. That result is not substantially different from the Canadian system, where private insurance pays for non-core services, but health care benefits are paid from general taxation revenues rather than by a mixture of public pay and work-related insurance policies.

The real test is whether the system promotes as part of its essential premise and structure a person's ability to "buy" his way to preferential care. This is not part of the basic framework on which medicare is based. Canada's single-payer system can continue to provide Canadians with quality care that is equitably accessible. The challenge for us is to use our resources creatively rather than to expand the private tier to include core clinical services within our health care system.<sup>5,6</sup>

### Michael Gordon, MD

Vice-President, Medical Services  
Head, Geriatric and Internal Medicine  
Baycrest Centre for Geriatric Care  
Head, Division of Geriatrics  
Department of Medicine  
Mount Sinai Hospital  
Professor of Medicine  
University of Toronto  
Toronto, Ont.

### References

1. Gray C. US editorial writers put Canadian health care under microscope. *CMAJ* 1998;159(11):1395-7.
2. Dirnfeld V. The benefits of privatization. *CMAJ* 1996;155:407-10.
3. Sky K. Should specialists support Canada's single-payer system? *Ann R Coll Physicians Surg Can* 1997;30:12-3.
4. Gunton RW. It's time to reintroduce a private component to the funding of health care. *Ann R Coll Physicians Surg Can* 1997;30:9-11.
5. Gordon M. A system worth saving. *CMAJ* 1996;154(9):1395-6.
6. Gordon M, Berger PB. The alluring myth of private medicine. *CMAJ* 1996;155(4):404-6.

## More than milk, eggs and orange juice

In his editorial on maternal nutrition,<sup>1</sup> Michael S. Kramer argues that the evidence does not strongly support the importance of maternal nutrition in determining the outcome of pregnancy in industrialized countries such as Canada. However, there

is clear evidence to support the importance of nutritional risk factors in the prevention of both broad subgroups of low birth weight: small-for-gestational-age births (which result from intrauterine growth retardation) and prematurity (which accounts for most low-birth-weight births in developed countries).

Low birth weight, which Kramer calls "*not* a very useful outcome" (emphasis in the original) has been described as "one of the most important biologic predictors of infant death and deficiencies in physical and mental development during childhood among those babies who survive."<sup>2</sup> The World Health Organization recommends the use of Williams' birth-weight curve to diagnose small-for-gestational-age births. The cut-off for a small-for-gestational-age term infant is 2900 g, and mounting evidence indicates that infants whose birth weight is above 2500 g but below about the 10th percentile still have higher health and nutrition risks than those whose birth weight is above the 10th percentile.<sup>3</sup>

In a 1987 meta-analysis<sup>3</sup> Kramer concluded that in developed countries the most important risk factors for intrauterine growth retardation were low maternal energy intake or weight gain and low pregravid weight, and that low pre-pregnancy weight was the only important nutrition-related risk factor for preterm birth. A more recent study found that both preterm labour and small-for-gestational-age births were associated with several factors, including low pre-pregnancy weight and low weekly maternal weight gain.<sup>4</sup> Other studies have found a relation between gestational weight gain and preterm delivery.<sup>5-7</sup> In 2 of these,<sup>6,7</sup> inadequate weight gain during the last half of pregnancy or the third trimester was associated with a higher risk of preterm birth.



Kramer attacks the Canada Prenatal Nutrition Program (CPNP) because it is based on the US Women, Infants, and Children (WIC) program. Yet on the basis of evidence that WIC reduced low-birth-weight births by 25% and very-low-birth-weight births by 44%, the US General Accounting Office concluded that WIC was a cost-effective program, resulting in savings of US\$2.89 to US\$3.50 for each federal dollar spent during the first 18 years of life.<sup>8</sup>

We should clarify that “providing milk, eggs and orange juice” is only one component of the CPNP, which addresses a number of issues that affect women’s overall health, including drug and alcohol use, smoking, family violence and social isolation, in addition to maternal nutrition and breastfeeding. We realize that Health Canada has a challenging task in evaluating the CPNP, and we look forward to the results. Although we agree with Kramer that more funding should be allocated to research, this should not occur at the expense of other worthwhile interventions.

**Sheela V. Basrur, MD, MHSc**

Medical Officer of Health

**Mary-Jo Makarchuk, MSc, MHSc, RD**

Public Health Nutritionist

Toronto Public Health

Toronto, Ont.

## References

1. Kramer MS. Maternal nutrition, pregnancy outcome and public health policy. *CMAJ* 1998;159(6):663-5.
2. Institute of Medicine. *WIC nutrition risk criteria: a scientific assessment*. Washington: National Academy Press; 1996.
3. Kramer MS. Determinants of low birth weight: methodological assessment and meta-analysis. *Bull World Health Organ* 1987;65(5):663-737.
4. Lang J, Lieberman E, Cohen A. A comparison of risk factors for preterm and term small-for-gestational-age birth. *Epidemiology* 1996;7:369-76.
5. Abrams B, Newman V. Small-for-gestational-age birth: maternal predictors and comparison with risk factors of spontaneous preterm delivery in the same cohort. *Am J Obstet Gynecol* 1991;164:785-90.
6. Abrams B, Newman V, Key T, Parker J. Maternal weight gain and preterm delivery. *Obstet Gynecol* 1989;74:577-83.
7. Siega-Riz AM, Adair LS, Hobel C. Maternal underweight and inadequate weight gain during the third trimester of pregnancy increases risk of preterm. *J Nutr* 1996;126:146-53.
8. Owen A, Owen G. Twenty years of WIC: a review of some effects of the program. *J Am Dietet Assoc* 1997;97:777-82.

## [The author responds:]

By continuing to espouse low birth weight as a useful outcome, Sheela Basrur and Mary-Jo Makarchuk persist in clouding the important distinction between preterm births and small-for-gestational-age births that perinatal epidemiologists and child health policy-makers have been trying to clarify over the last 15 to 20 years. Preterm birth, particularly birth before 32 completed weeks of gestation, is indeed “one of the most important biologic predictors of infant death and deficiencies in physical and mental development during childhood among those babies who survive.” However, small-for-gestational-age birth is not. And the evidence is quite clear that maternal nutrition in pregnancy has a far greater impact on small-for-gestational-age birth than on preterm birth.

Basrur and Makarchuk have been selective in citing references that support their prior belief and have failed to orient the reader to the methodologic strengths and weaknesses of the studies cited. None of the “positive” studies mentioned was a randomized trial. Basrur and Makarchuk argue that nutrition is just one component of the CPNP, but randomized trials of routine advice to reduce smoking<sup>1</sup> or of providing psychosocial support<sup>2</sup> or intensive prenatal care<sup>3-5</sup> to high-risk women have yielded consistently negative results. Even the recently published trial of balanced energy-protein supplementation in marginally nourished pregnant women in a rural area of The Gambia, which reported large beneficial effects on fetal growth (i.e., birth

weight for gestational age), found no effect whatsoever on the duration of gestation.<sup>6</sup> Surely the results of systematic reviews and large individual trials should take precedence over selectively cited observational studies.

Proponents of the WIC program in the US cite evidence of effectiveness from comparisons of WIC participants and nonparticipants. But women who participate in WIC (or any other public health program, for that matter) are different from those who do not. Women who deliver very early, for example, will not have had the same time to enrol in WIC as those who deliver at term. Thus preterm birth can lead to nonparticipation, and an observational study may well put the cart before the horse by attributing to WIC the lower rate of preterm birth among participants. Participants also tend to be more committed to their pregnancies, are more conscious of their health in general, have the psychological and financial wherewithal to enrol and attend WIC clinic visits and are likely to eat better on their own. It is quite impossible to control for such potent confounding effects. Because no amount of replication using a similar scientifically flawed design can replace comparison based on randomized allocation, the “evaluation” of CPNP so eagerly awaited by Basrur and Makarchuk will be as useless as the WIC evaluations.

I am also concerned about the belief that randomized trials are splendid tools for evaluating health care interventions in individuals but are unfeasible or unethical for evaluating community interventions. To be sure, randomization of individuals living in the same community is difficult because of the inevitable dilutional effect (“contamination”) caused by shared experiences and behaviours. Is it for this reason that cluster randomization (in which the clinic, the hospital or the entire community becomes the unit of randomization) has be-



come such a powerful tool. Examples of perinatal trials using cluster randomization include studies of counts of fetal movement in the prevention of antepartum stillbirth,<sup>7</sup> of early breast-feeding to prevent postpartum hemorrhage<sup>8</sup> and of counselling for smoking cessation in prenatal care,<sup>1</sup> as well as a WHO-sponsored evaluation of a new model of prenatal care.<sup>9</sup> My colleagues and I are currently conducting a cluster-randomized evaluation (funded in part by Health Canada) of an intervention to promote breast-feeding based on the WHO/UNICEF Baby-Friendly Hospital Initiative. Cluster-randomized trials require highly trained research teams, large sample sizes and substantial funding. If individual-based interventions deserve rigorous methods of evaluation, the far larger number of individuals whose health and welfare may be affected argues for better, not worse, methods of evaluating community-based interventions.

I have no objection to funding truly "worthwhile interventions," whose effectiveness has been rigorously demonstrated. In the maternal-child health arena alone, postnatal support of breast-feeding, provision of automobile restraints and bicycle helmets, and improvement in vaccination coverage are public health promotion efforts whose scientific basis is far stronger than that of CPNP.

#### Michael S. Kramer, MD

Professor  
Departments of Pediatrics  
and of Epidemiology and Biostatistics  
Faculty of Medicine  
McGill University  
Montreal, Que.

#### References

1. Kendrick JS, Zahniser SC, Miller N, Salas N, Stine J, Gargiullo PM, et al. Integrating smoking cessation into routine public prenatal care: the Smoking Cessation in Pregnancy Project. *Am J Public Health* 1995;85:217-22.
2. Hodnett ED. Support from caregivers

during at-risk pregnancy. In: Neilson JP, Crowther CA, Hodnett ED, Hofmeyr GJ, editors. *Pregnancy and childbirth module of the cochrane database of systematic reviews* [updated 1997 Dec 2]. Available: The Cochrane Library [database on disk and CD-ROM]. The Cochrane Collaboration; Issue 1. Oxford: Update Software; 1998.

3. Mueller-Heubach E, Reddick D, Barnett B, Bente R. Preterm birth prevention: evaluation of a prospective controlled randomized trial. *Am J Obstet Gynecol* 1989; 160:1172-8.
4. Heins HC, Nance NW, McCarthy BJ, Efrid CM. A randomized trial of nurse-midwifery prenatal care to reduce low birth weight. *Obstet Gynecol* 1990;75:341-5.
5. Collaborative Group on Preterm Birth Prevention. Multicenter randomized, controlled trial of a preterm birth prevention program. *Am J Obstet Gynecol* 1993;169: 352-66.
6. Ceesay SM, Prentice AM, Cole TJ, Foord F, Weaver LT, Poskitt EME, et al. Effects on birth weight and perinatal mortality of maternal dietary supplements in rural Gambia: 5 year randomised controlled trial. *BMJ* 1997;315:786-90.
7. Grant A, Elbourne D, Valentin L, Alexander S. Routine formal fetal movement counting and risk of antepartum late death in normally formed singletons. *Lancet* 1989;2:345-9.
8. Bullough C, Msuku R, Karonde L. Early suckling and postpartum hemorrhage: controlled trial in deliveries by traditional birth attendants. *Lancet* 1989;2:522-5.
9. Villar J, Bakketeig L, Donner A, al-Mazrou Y, Ba'aqeel H, Belizán JM, et al. The WHO Antenatal Care Randomised Controlled Trial: rationale and study design. *Paediatr Perinat Epidemiol* 1998;12 (Suppl 2):27-58.

### What causes chronic fatigue?

Even though the 3 articles on chronic fatigue syndrome<sup>1-3</sup> in the Sept. 8 issue commendably demolish the obsolete claim that chronic fatigue syndrome is a psychiatric illness, they also offer outdated biological explanations for the syndrome, namely, either a chronic viral infection or a weakened immune system. Although the first of these explanations seemed convincing until a few years ago, it is hardly tenable now, because no specific virus has been identified in these patients.<sup>4</sup>

Both the viral reactivation and the immunological abnormalities observed in patients with chronic fa-

tigue syndrome may well be accounted for by the cortisol deficiency that characterizes these patients.<sup>5</sup> This explanation is supported by the striking similarities between chronic fatigue syndrome and Addison's disease, which share 26 features,<sup>6</sup> including all of the neuropsychological symptoms.<sup>5</sup>

My conviction that chronic fatigue syndrome is an adrenal insufficiency similar to Addison's disease lies primarily in the fact that 4 years ago I recovered from chronic fatigue syndrome in the course of a few days thanks to the consumption of licorice,<sup>7</sup> with which addisonian patients were successfully treated before hydrocortisone and fludrocortisone became available.<sup>7</sup> These steroids, which currently represent the lifelong therapy for Addison's disease,<sup>7</sup> should be investigated in the treatment of patients with "true" chronic fatigue syndrome, as diagnosed according to the original criteria.<sup>8</sup> Conversely, patients in whom chronic fatigue syndrome is diagnosed on the basis of subsequent revised criteria<sup>9</sup> (which do not include the only physical signs — enlarged lymph nodes, fever and sore throat — that clearly distinguish chronic fatigue syndrome from depression) should avoid both steroid replacement therapy and licorice. In fact, depressed patients misdiagnosed as having chronic fatigue syndrome have abnormally high cortisol levels,<sup>10</sup> instead of the abnormally low cortisol levels found in patients with "true" chronic fatigue syndrome.<sup>10</sup> Therefore, administration of licorice or hydrocortisone would further increase their already-high cortisol levels.<sup>7</sup>

#### Riccardo Baschetti, MD

Padua, Italy

#### References

1. Caplan C. Chronic fatigue syndrome or just plain tired? *CMAJ* 1998;159(5):519-20.
2. Capen K. Chronic fatigue syndrome gets court's nod of approval as legitimate disorder. *CMAJ* 1998;159(5):533-4.
3. Sibbald B. Chronic fatigue syndrome