

Challenges in family practice related to informed and shared decision-making: a survey of preceptors of medical students

William Godolphin, Angela Towle, Rachael McKendry

§ See related article page 438

Although the informed involvement of patients in shared decision-making is commonly acknowledged to be an ideal situation, this is rarely achieved,¹ leading to important adverse consequences.² It can be a communication challenge for clinicians to involve patients in informed and shared decision-making (ISDM). We defined the “competencies” needed for ISDM³ in order to address this problem systematically through medical education. These competencies are communication skills related to building of partnerships; eliciting of preferences for receiving information and playing a role in decision-making; exploring of ideas, concerns and expectations; presenting of choices and evidence; reaching a decision and resolving conflict; and agreeing on an action plan and follow-up.

In this study, we asked, What are the most frequent and challenging situations that require ISDM skills experienced by family practice preceptors of medical students? The information could be used to refine training in communication skills, because students and preceptors are likely to be motivated to learn approaches to difficult and common problems.

It has been reported that the “difficult patient” occurs in 10%–20% of primary care encounters,⁴ but the terms used in the substantial “difficult patient” literature do not seem to be congruent with the “difficulties” that physicians may have in engaging patients in ISDM. Some examples that do seem relevant have been suggested by Platt and Gordon:⁵ “nonadherence,” “the list maker,” “the patient’s companion,” “the patient bearing literature” and, in a survey of general practitioners’ frequent problems, “patient’s non-compliance with treatment and follow-up,” “sharing understanding of the problem with the patient” and “differences in expectations between physician and patient.”⁶

We wrote survey questions to portray generic situations in physician–patient encounters that pose a challenge to ISDM. The scenarios were developed and refined by step-wise and iterative consultation with physicians who had wide experience in teaching and examining communication skills and had helped to define the competencies for ISDM. The survey form was constructed with the attention to de-

Box 1: Survey questions used to measure the perceived frequency and challenges of physician–patient encounters that may be addressed with informed and shared decision-making (ISDM) skills

How challenging are the following situations for you in your practice and how often are you faced with them?

1. Dealing with a patient who does not respond to treatment as expected and admits to not taking medication as prescribed (NonComply)
2. Responding to a patient who wants to try an alternative or complementary therapy about which you have major concerns (A/CTherapy)
3. Dealing with a patient who wants something (e.g., a test, prescription or referral) that you do not think is appropriate or necessary (RxConflict)
4. Responding to a patient who has a lot of information (e.g., from books, the Internet, friends) but is unable to assess its quality (LotsOfInfo)
5. Handling a situation in which the patient is accompanied by a significant other (e.g., spouse, parent) who interferes with your ability to identify the problem or discuss treatment (SOInterfere)
6. Conducting an interview in which you suspect that the patient is shopping for a doctor (ShopForDoc)
7. Managing a patient who has a progressive chronic condition and refuses the best management option despite being presented with the evidence (RefuseBest)
8. Responding to a patient who wants to know your opinion about a therapy that you do not know about (conventional or alternative/complementary therapy) (DrNotKnow)
9. Involving a patient in making a choice between different treatments (or investigations) when they want you to make the decision and you feel that this is not appropriate (YouDecide)

tail and format suggested by Woodward.⁷ Problems were expressed as 9 items, for example, “dealing with a patient who does not respond to treatment as expected and admits to not taking medication as prescribed” (Box 1). Respondents were asked to what degree they felt challenged by the problem and how frequently they encountered it on 5-point scales from “1: not challenging (you feel effective and easily able to address the problem and it causes you little anxiety)” to “3: moderately challenging” to “5: very challenging (you find this to be quite a difficult problem and anxiety provoking)” and from “1: once a year” to “3: once a month” to “5: most days.” An illustrative example (dealing with a patient who is “drug-seeking”) was used at the beginning of the questionnaire as an anchor for the scale.

The survey was mailed to all 285 family practice preceptors of medical students at the University of British Columbia. The survey was anonymous. Guidelines for maximizing returns were followed.⁸ The response rate was 88% and the survey showed high internal consistency, with a Cronbach α of 0.82 for challenge and 0.84 for frequency. Of the physicians who completed the survey, 51% were local and 49% were rural; 66% were in group practices, 26% in solo practices, 1% in walk-in clinics and 7% indicated “other”;

and 74% were male. The average time in practice was 14 years, with a range from 0.5 to 42 years.

The most challenging problems were conflict resolution, dealing with “significant others” and patients’ refusal of best treatment. The most frequent problems were conflict resolution, noncompliance and patients with lots of information (Fig. 1). There were no important correlations with physicians’ type of practice, sex or years of experience.

Our results indicate that training in communication skills for undergraduates should include attention to conflict resolution and negotiation skills and that their preceptors also need help with this. Training should also provide students with strategies for managing decision-making in the context of patients and their companions.

This article has been peer reviewed.

Dr. Godolphin is Professor, Department of Pathology and Laboratory Medicine, and Codirector, Faculty Development, Faculty of Medicine, University of British Columbia, Vancouver, BC. Dr. Towle is Director, MD Undergraduate Program, Faculty of Medicine, University of British Columbia, Vancouver, BC. Ms. McKendry is Research Assistant, Informed Shared Decision Making Project, Office of the Coordinator of Health Sciences, University of British Columbia, Vancouver, BC.

Competing interests: None declared.

Contributors: The conceptualization of the study was primarily the work of Drs. Towle and Godolphin. All 3 authors contributed to interviews with physicians to define the questionnaire items and to the detailed design of the study. The mailing of the survey, data collection and statistical data analysis were mainly carried out by Ms. McKendry, who also wrote a first draft of the paper. The final version of the manuscript was written mainly by Drs. Godolphin and Towle.

Acknowledgements: We are especially indebted to the physicians who helped us create and carry out the survey: Drs. Stephen Barron, Betty Calam, Stefan Grzybowski, Jacqueline Hurst, Joy Masuhara, Katherine McManus, James Salzman, Robert Voigt, Arthur van Wart, Robert Woollard (Department of Family Practice) and Andrew Chalmers (Department of Medicine).

The protocol was approved by the University of British Columbia Behavioural Research Ethics Board. This work was supported by a grant from The Max Bell Foundation.




Fig. 1: Percentage of responses from family practice preceptors for level of frequency (top) and challenge (bottom) for each item in the questionnaire. For expansions of abbreviations, see Box 1.

References

1. Braddock CH, Edwards KA, Hasenberg NM, Laidley TL, Levinson W. Informed decision making in outpatient practice. Time to get back to basics. *JAMA* 1999;282:2313-20.
2. Britten N, Stevenson FA, Barry CA, Barber N, Bradley CP. Misunderstandings in prescribing decisions in general practice: qualitative study. *BMJ* 2000; 320:484-8.
3. Towle A, Godolphin W. Framework for teaching and learning informed shared decision making. *BMJ* 1999;319:766-71.
4. Hahn SR, Kroenke K, Spitzer RL, Brody D, Williams JBW, Linzer M, et al. The difficult patient: prevalence, psychopathology, and functional impairment. *J Gen Intern Med* 1996;11:1-8.
5. Platt FW, Gordon GH. *Field guide to the difficult patient interview*. Philadelphia: Lippincott Williams & Wilkins; 1999.
6. Leclère H, Beaulieu MD, Bordage G, Sinden A, Couillard M. Why are clinical problems difficult? General practitioners' opinions concerning 24 clinical problems. *CMAJ* 1990;143(12):1305-15.
7. Woodward CA. Questionnaire construction and question writing for research in medical education. *Med Educ* 1988;22:347-63.
8. Dillman DA. *Mail and telephone surveys: the total design method*. New York: John Wiley & Sons Inc; 1978.

Correspondence to: Dr. William Godolphin, Rm. G227, 2211 Wesbrook Mall, University of British Columbia, Vancouver BC V6T 2B5; fax 604 822-7635; wgod@unix.ubc.ca