

- torial]. *CMAJ* 2004;170(10):1567-8.
- Lindstrom J. Isotretinoin: background and regulatory history [slide presentation]. In: Joint session of Drug Safety and Risk Management Advisory Committee and Dermatologic and Ophthalmic Drugs Advisory Committee; 26 Feb 2004; Gaithersburg (MD). Rockville (MD): US Food and Drug Administration; 2004. Available: [www.fda.gov/ohrms/dockets/ac/04/slides/4017S1\\_01\\_Linstrom.ppt](http://www.fda.gov/ohrms/dockets/ac/04/slides/4017S1_01_Linstrom.ppt) (accessed 2004 Nov 16). Slide 27: Number of reported pregnancies 1982-1999 by year of therapy (US).
  - Mitchell AA, Van Bennekom CM. Isotretinoin survey [slide presentation]. In: Joint session of Drug Safety and Risk Management Advisory Committee and Dermatologic and Ophthalmic Drugs Advisory Committee; 26 Feb 2004; Gaithersburg (MD). Rockville (MD): US Food and Drug Administration; 2004. Available: [www.fda.gov/ohrms/dockets/ac/04/slides/4017S1\\_04\\_Mitchell.ppt](http://www.fda.gov/ohrms/dockets/ac/04/slides/4017S1_04_Mitchell.ppt) (accessed 2004 Nov 16). Slide 4: Pregnancy rate/1000 person courses by enrolment year.

*Competing interests:* None declared. The views expressed are those of the authors and do not necessarily reflect those of Health Canada.

DOI:10.1503/cmaj.1041643

### [Two of the authors respond:]

As outlined in our commentary,<sup>1</sup> we agree that every effort should be made to prevent fetal exposure to isotretinoin, a highly teratogenic drug.

Maria Valois and associates state that, unlike the situation in the United States, Health Canada has no evidence of a significant rate of pregnancy among Canadian women taking isotretinoin. Unfortunately, this impression is incorrect. The presentations to the FDA in February 2004, attended by Health Canada, included a study by the Organization of Teratology Information Services,<sup>2</sup> which was funded by the US Centers for Disease Control and Prevention. More than half of the cases in that study came from Canada. The Canadian data were collected prospectively in 2002-2003 by The Motherisk Program (based in Toronto) and IM-AGE (based in Montréal). On a proportional basis, Canada had substantially more cases than the United States.

This situation reflects the understandable ineffectiveness of reporting systems based on spontaneous reports. Moreover, in 2003 Health Canada stopped the development of MotherNet, a system that would have given the department such information continuously. The reason cited for the halt was lack of funding, although over \$1 million had been spent on the project at that point.

Wouldn't it make sense for Health Canada to work with existing active surveillance programs in Toronto and Montréal and thus to receive the tremendous amount of data that are being collected on drugs in pregnancy in Canada? Presently, no arrangements exist for such collaboration.

#### Gideon Koren

Director, The Motherisk Program

#### Neil Shear

Helen and Paul Phalen Professor

of Dermatology

Chief of Dermatology

University of Toronto

Toronto, Ont.

#### References

- Koren G, Avner M, Shear N. Generic isotretinoin: a new risk for unborn children [editorial]. *CMAJ* 2004;170(10):1567-8.
- Interim report OTIS — North American isotretinoin information and survey line [slide presentation]. In: Joint session of Drug Safety and Risk Management Advisory Committee and Dermatologic and Ophthalmic Drugs Advisory Committee; 26 Feb 2004; Gaithersburg (MD). Rockville (MD): US Food and Drug Administration; 2004. Available: [www.fda.gov/ohrms/dockets/ac/04/slides/4017S1\\_09\\_Miller-OTIS.ppt](http://www.fda.gov/ohrms/dockets/ac/04/slides/4017S1_09_Miller-OTIS.ppt) (accessed 2004 Nov 16).

*Competing interests:* Neil Shear has received educational grants from Hoffmann-La Roche.

DOI:10.1503/cmaj.1041700

### Outcomes of postmastectomy radiotherapy

In 1995 the American Society of Clinical Oncology adopted a statement on the choice of outcomes in assessing cancer treatments.<sup>1</sup> That statement made a clear distinction between patient outcomes (survival and quality of life) and cancer outcomes (tumour regression), the former being much more important. This view is also expressed in current books on cancer therapy.<sup>2</sup> A recently published guideline, though devoted to techniques of measuring tumour response, stated that this outcome is of value as an endpoint in early clinical trials, but in phase III trials and clinical application "it should not be the sole, or major, endpoint."<sup>3</sup> Yet the clinical practice guidelines published in *CMAJ* concerning the use of postmastectomy radiotherapy<sup>4</sup> appear to be founded entirely on evidence related to

tumour responsiveness. Although local irradiation is effective in destroying local tumour tissue, none of the relevant clinical trials have shown that this leads to an improvement in overall survival. Normally, acceptance of a therapeutic modality requires demonstration of its efficacy, yet leading oncologists appear to take the opposite stance in regard to radiotherapy. Thus it is assumed, despite a lack of supporting evidence, that the majority of patients "require" irradiation but that subgroups who do not benefit will ultimately become recognizable.<sup>5,6</sup>

The authors of the guidelines are to be commended for including a "questions and answers" guide for women and their physicians (Appendix 1 of the article).<sup>4</sup> But one important question has been omitted: "How will radiation help me?" One must wonder how many of the patients anxiously waiting for radiation therapy are among those for whom therapy has been recommended despite a lack of evidence of benefit. If such patients were given balanced information and allowed to choose whether to undergo therapy, how many would decide against it?

#### Norman Kalant

Sir Mortimer B. Davis Jewish

General Hospital

Montréal, Que.

#### References

- American Society of Clinical Oncology. Outcomes of cancer treatment for technology assessment and cancer treatment guidelines. *J Clin Oncol* 1996;14:671-9.
- Weeks JC. Outcomes assessment. In: Kufe DW, editor. *Holland-Frei cancer medicine*. 6th ed. Hamilton (ON): BC Decker; 2003. p. 479-84.
- Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, et al. New guidelines to evaluate the response to treatment in solid tumors. *J Natl Cancer Inst* 2000;92:205-16.
- Truong PT, Olivetto IA, Whelan TJ, Levine M, for the Steering Committee on Clinical Practice Guidelines for the Care and Treatment of Breast Cancer. Clinical practice guidelines for the care and treatment of breast cancer: 16. Locoregional post-mastectomy radiotherapy. *CMAJ* 2004;170(8):1263-73.
- Winer EP, Morrow M, Osborne CK, Harris JR. Malignant tumors of the breast. In: DeVita VT Jr, Hellman S, Rosenberg SA, editors. *Cancer: principles and practice of oncology*. 6th ed. Philadelphia: Lippincott Williams & Wilkins; 2001. p. 1651-717.
- Margolese RG, Hortobagyi GN, Buchholz TA. Neoplasms of the breast. In: Kufe DW, Pollock RE, Weichselbaum RR, Bast RC Jr, Gansler TS, Holland JF, et al, editors. *Holland-Frei cancer medicine*. 6th ed. Hamilton (ON): BC Decker; 2003. p. 1879-972.

DOI:10.1503/cmaj.1041280