

# A clinical return-to-work rule for patients with back pain

Clermont E. Dionne, Renée Bourbonnais, Pierre Frémont, Michel Rossignol, Susan R. Stock, Isabelle Larocque

β See related article page 1575

## Abstract

**Background:** Tools for early identification of workers with back pain who are at high risk of adverse occupational outcome would help concentrate clinical attention on the patients who need it most, while helping reduce unnecessary interventions (and costs) among the others. This study was conducted to develop and validate clinical rules to predict the 2-year work disability status of people consulting for nonspecific back pain in primary care settings.

**Methods:** This was a 2-year prospective cohort study conducted in 7 primary care settings in the Quebec City area. The study enrolled 1007 workers (participation, 68.4% of potential participants expected to be eligible) aged 18–64 years who consulted for nonspecific back pain associated with at least 1 day's absence from work. The majority (86%) completed 5 telephone interviews documenting a large array of variables. Clinical information was abstracted from the medical files. The outcome measure was "return to work in good health" at 2 years, a variable that combined patients' occupational status, functional limitations and recurrences of work absence. Predictive models of 2-year outcome were developed with a recursive partitioning approach on a 40% random sample of our study subjects, then validated on the rest.

**Results:** The best predictive model included 7 baseline variables (patient's recovery expectations, radiating pain, previous back surgery, pain intensity, frequent change of position because of back pain, irritability and bad temper, and difficulty sleeping) and was particularly efficient at identifying patients with no adverse occupational outcome (negative predictive value 78%–94%).

**Interpretation:** A clinical prediction rule accurately identified a large proportion of workers with back pain consulting in a primary care setting who were at a low risk of an adverse occupational outcome.

Intervention for back pain has great potential for improving population health and for freeing up extensive societal resources.

So-called red flags to identify pain that is specific (i.e., pain in the back originating from tumours, fractures, infections, cauda equina syndrome, visceral pain and systemic disease)<sup>5</sup> account for about 3% of all cases of back pain.<sup>6</sup> The overwhelming majority of back-pain problems are thus nonspecific. One important feature of nonspecific back pain among workers is that a small proportion of cases (< 10%) accounts for most of the costs (> 70%).<sup>7–14</sup> This fact has led investigators to focus on the early identification of patients who are at higher risk of disability, so that specialized interventions can be provided earlier, whereas other patients can be expected to recover with conservative care.<sup>9,15–25</sup> Although this goal has become much sought-after in back-pain research, most available studies in this area have 3 methodological problems:

- Potential predictors are often limited to administrative or clinical data, whereas it is clear that back pain is a multidimensional health problem.
- The outcome variable is most often a 1-point dichotomous measure of return to work, time off work or duration of compensation, although some authors have warned against the use of first return to work as a measure of recovery. Baldwin and colleagues,<sup>26</sup> for instance, point out that first return to work is frequently followed by recurrences of work absence.
- Most published prediction rules developed for back pain have not been successfully validated on any additional samples of patients.

Our study aimed to build a simple predictive tool that could be used by primary care physicians to identify workers with nonspecific back pain who are at higher risk of long-term adverse occupational outcomes, and then to validate this tool on a fresh sample of subjects.

## Methods

From 1998 through 2002, we conducted a prospective cohort study called the RAMS-Prognosis Study (RAMS stands for *recherche sur les affections musculo-squelettiques*), which had a qualita-

CMAJ 2005;172(12):1559-67

Since the 1950s, back pain has taken on the proportions of a veritable epidemic, counting now among the 5 most frequent reasons for visits to physicians' offices in North America<sup>1–3</sup> and ranking sixth among health problems generating the highest direct medical costs.<sup>4</sup> Because of its high incidence and associated expense, effec-

tive and a quantitative phase. The study was approved annually by the ethics committees of the participating hospitals.

In the first phase, 2 focus groups composed of people who have had severe back pain were held in November and December, 1998. The focus groups enriched the list of potential predictors (Appendix 1)<sup>10,27</sup> to be investigated quantitatively in the interview-based second phase. The variables they identified were added to a list of those collected from the existing literature, to document as many potential predictors as possible.

Subjects for the second phase were recruited in 7 primary care settings of the Quebec City area: 4 emergency departments and 3 family medicine units, all having medical teaching responsibilities. Subjects of interest were adult workers aged 18–64 years who consulted for back pain, whatever its character or duration, between June 1999 and September 2000. Patients were eligible if the back pain was nonspecific and had caused them to be absent from their regular job for at least 1 day. Excluded were patients with any other condition that could affect their work capacity (pregnancy, for example, or a serious comorbidity) and those whose pain was located in the cervical spine only or had a specific cause.

Lists of potential subjects at each site were submitted weekly. A medical archivist telephoned these patients to request their participation in the study; those who agreed and were eligible were mailed an informed consent form to sign and return.

**Table 1: Definitions of “return to work in good health” outcome categories**

Category	Definition
Success	At regular work (i.e., like that performed before the index medical consultation for back pain) as of day of interview and Functional limitations $\leq 30\%^*$ and No more than 5 days of absence from regular work since last interview ( $\leq 7$ d, for the interviews at 1 and 2 yr) OR Other situation unrelated to back pain (e.g., became homemaker by choice)
Partial success	
PS1	Attending regular work, as of day of interview and Functional limitations $\leq 30\%$ and More than 5 days of absence from regular work since last interview ( $> 7$ d, for the interviews at 1 and 2 yr) OR
PS2	Attending regular work and Functional limitations $> 30\%$ and No more than 5 days of absence from regular work since last interview ( $\leq 7$ d, for the interviews at 1 and 2 yr) OR
PS3	Attending regular work and Functional limitations $> 30\%$ and More than 5 days of absence from regular work since last interview ( $> 7$ d, for the interviews at 1 and 2 yr)
Failure after attempt	Ongoing absence from regular work because of back pain, as of day of interview and At least 1 attempt to return to regular work since last interview
Failure	Still absent from regular work because of back pain and No attempt to return to work as yet

\*Functional limitations were measured with a Canadian French translation of the Roland–Morris Disability Questionnaire.

Participants were telephoned by trained interviewers for a baseline interview at around 3 weeks after their medical consultation, with repeated measurements at 6 and 12 weeks and 1 and 2 years. (Note that all timeframes are relative to each patient’s index medical consultation.) The data collected described demographic, socioeconomic, behavioural, anthropometric, clinical, occupational and psychosocial variables, as well as information on the utilization of health services for back pain. Clinical information was drawn by a single individual from study participants’ medical files. An online appendix (available at [www.cmaj.ca/cgi/content/full/172/12/1559/DC1](http://www.cmaj.ca/cgi/content/full/172/12/1559/DC1)) summarizes the standardized instruments used to measure key constructs.<sup>28–60</sup> When an instrument was unavailable in French, Vallerand’s double-inverse translation method<sup>61</sup> was applied.

Return to work in good health (RWGH) is an index of back-pain outcome that takes into account work status, functional limitations and number of days of work-absence since the medical consultation. This variable has 4 categories based on the work of Baldwin’s group<sup>26</sup> (presented in detail in Table 1): success, partial success, failure after attempt, and failure. Our work-absence data were self-reported; however, a validation study was conducted with data obtained from the employers of 40 volunteers. The type 1 intraclass coefficient correlating the 2 sets of data was 0.97 (95% confidence interval [CI] 0.93–0.98) for number of days of modified work, and 0.99 (95% CI 0.98–1.00) for number of days off any work.

Subsequent to descriptive analysis, participants were randomly assigned (by means of a randomization function in SAS software) into 2 subgroups: about 40% into a “training sample” and the rest into a “validation sample.” Predictive analyses were conducted to identify, not causal relationships, but only prognostic indicators that could be clinically useful.<sup>62</sup> Recursive partitioning with KnowledgeSeeker software (version 3.0, Angoss Software International Ltd., Toronto, Ont.) was used to build a predictive model of 2-year RWGH in the training sample.<sup>63</sup> Because the distinction between the “failure after attempt” and “failure” groups has no clinical pertinence, data for these 2 categories of outcome were lumped together. The threshold for statistical significance was fixed first at 0.01, to favour the selection of the strongest predictors and simplicity in the model built. Afterward, it was relaxed to  $p \leq 0.05$  to permit us to identify complementary and alternative associations.

Classification error rate, sensitivity, specificity, and positive and negative predictive values (the probability, respectively, of having and not having an actual adverse occupational outcome when so predicted by the model) were computed for each model by comparing the predicted classification of subjects to their actual status at 2 years. In these analyses, the subjects were dichotomized along different groupings of the outcome categories. The “success” category was always considered among the “non-diseased” group. Area under the receiver operating characteristic curve<sup>64</sup> was used as the main criterion to select one model over another, followed by the number of variables. Each selected model was then applied to the validation sample. We “pruned” the final model to try to keep it as simple as possible. All baseline, 6-week and 12-week variables were candidate predictors, including individual items or questions of specific measurement tools (Appendix 1).


## Results

Numbers of eligible subjects, refusals and participants are shown in Fig. 1. The proportion of eligibility among subjects who were reached (37.6%) was applied to those

not reached, to estimate a total number of eligible subjects. A total of 1007 subjects (68.4% of the 1471 expected to have been eligible) participated in the baseline interview, conducted an average of 25 days after the index consultation (standard deviation [SD] 10.2 d). Of those who participated in the baseline interview, 923 (91.7%) completed the interview at 6 weeks of follow-up, 907 (90.1%) at 12 weeks, 913 (90.7%) at 1 year, and 864 (85.8%) at 2 years. Three patients died during the follow-up period. Completed records of all 5 interviews were available for 860 participants, whose data were used for the recursive partitioning analysis.

The mean age of the participants was just under 39 years; a majority (58.5%) were male (Table 2). Almost half (47.8%) had earned a postsecondary diploma, from either a community college or a university.

At baseline, a majority of subjects reported their back pain to be recurrent (“it comes and goes”) or persistent (“the pain is always there, to different degrees”); fewer than one-quarter reported that theirs was a one-time problem (“never had back pain before”). The median time since the beginning of patients’ first episode was nevertheless 6 years. Pain was mostly situated in the lumbar and lumbosacral



**Fig. 1: Eligible subjects, refusals and participants in the study.** \*Eligibility unknown. R = randomization.

**Table 2: Selected sociodemographic variables of study participants (workers with back pain) at the baseline interview**

Characteristic*	No. (%), n = 1007	Mean (SD)	Data missing
Age, yr (median 39.0, range 18–64)		38.7 (10.6)	0
Female	418 (41.5)		0
Graduated from primary school	186 (18.9)		24
Secondary school	327 (33.3)		
Community college	305 (31.0)		
University	165 (16.8)		
Body mass index < 27 kg/m <sup>2</sup>	682 (69.6)		27
≥ 27 kg/m <sup>2</sup>	298 (30.4)		
Currently smokes cigarettes	388 (39.2)		17
Former smoker	282 (28.5)		
Never smoked	320 (32.3)		
Duration of current back pain, d (median 4.0, range 1–8760 d)		70.8 (489.6)	3
Type of back pain: one-time	222 (22.2)		5
Recurrent; comes and goes	458 (45.7)		
Persistent; always present	322 (32.1)		
Pain is cervicothoracic	17 (1.8)		54
Thoracic	166 (17.2)		
Thoracolumbar	69 (7.2)		
Lumbar	533 (56.2)		
Lumbosacral or sacral	161 (16.9)		
At multiple sites	7 (0.7)		
Pain radiating to arms or legs	520 (56.4)		85
No pain radiation	402 (43.6)		
Period since first back pain, yr (median 6.0, range 0.003–45.1 yr)		8.7 (8.9)	4
Duration (patient-reported) of index medical consultation, min (range 1–240 min)		16.0 (14.9)	22
No bed rest recommended by MD	812 (84.2)		43
1–3 days recommended	61 (6.3)		
> 3 days recommended	91 (9.4)		
Since index medical consultation:			16
Completely satisfied with health services received	370 (37.3)		
Mostly satisfied	434 (43.8)		
Mostly unsatisfied	106 (10.7)		
Not at all satisfied	81 (8.2)		
Compensated for previous back pain	294 (29.8)		19
No previous compensation	694 (70.2)		
Unionized job	450 (45.5)		18
Not unionized	539 (54.5)		
Job seniority, yr		7.5 (8.3)	19
Index (frequency × effort) of job’s physical demands (MR 0–44)		18.1 (12.2)	21
No job satisfaction	38 (3.9)		16
Very low satisfaction	47 (4.7)		
Moderate satisfaction	341 (34.4)		
Complete satisfaction	565 (57.0)		
Symptoms of depression (MR 0–4)		1.5 (0.9)	21
Somatization symptoms (MR 0–4)		1.1 (0.8)	21
Self-efficacy (MR 0–100)		49.5 (23.3)	89

Note: SD = standard deviation, MR = maximum range of scores possible.  
\*Medians are provided when the data were not normally distributed (not symmetrical).

areas. Over half of our study subjects reported pain radiating to the arms or legs.

The 2-year evolution of RWGH among study subjects is illustrated in Fig. 2. The most important changes in RWGH occurred at about 12 weeks, at which time about 50% of subjects were in the RWGH success category (compared with 18% at 6 weeks). At 2 years, close to 20% were still in the “failure after attempt(s)” and “failure” groups.

Fig. 3 presents the final predictive model as a clinical algorithm that allows estimation of the probability, for a given individual, of RWGH success, partial success and failure (which includes failure after attempt). The classification error rate was 37.0% in the training sample and 40.5% in the validation sample. Measures of validity to detect RWGH failure, partial success or both outcomes taken together for this model are presented for both samples in Table 3, with a set of example calculations shown in Fig. 4. All validity measures were quite stable when applied to the validation sample. In all cases, findings for negative predictive value were high (74%–91%). It was highest for predicting “failure after attempt/failure” (91%), whereas the highest positive predictive value was for detecting the combined outcome of “failure after attempt/failure” or “partial success” simultaneously (57%).

## Interpretation

This study corroborates the complex nature of back pain and the inherent difficulty in developing clinical prediction tools for such conditions. With a large coverage of potential predictors measured at baseline and at 6 and 12 weeks, the best model we obtained contains 7 predictors measured at baseline (patient’s recovery expectations, radiating pain, previous back surgery, intense pain, frequent


change of position because of back pain, irritability and bad temper, and difficulty sleeping). It is far from perfect; nonetheless, its high negative predictive value may constitute a strong advantage.

Most previous studies on the prediction of the long-term outcome of back pain used a dichotomous measure of return to work, duration of work absence or compensation data. Because of methodological differences and the nature of predictive analyses (which are not intended to identify causal relationships, and should not be interpreted as doing so),<sup>64</sup> it is not relevant to directly compare the predictors. In fact, 2 studies could end up with different sets of predictors; but the nature of the predictors is not very important, as long as they constitute an efficient and reproducible prognostic tool. The comparison must be made on the predictive validity of the models, that is, on their capacity to classify subjects correctly with respect to their outcome and on demonstration of the reproducibility of this validity. The few existing previously determined models explained some 25%–30% of the variance of continuous outcomes,<sup>25,65</sup> similar to what we observed when we applied our final predictive model to long-term absence (data not shown). With regard to other measures of predictive validity, it is quite often negative predictive values that are highest,<sup>65-67</sup> as in our study.


This study had several strengths: the inclusion of all workers consulting for back pain, whatever the source of pain (i.e., not only workers’ compensation cases); the large sample; the prospective design, with repeated measures taken at key points in the natural history of the disease; a high participation rate; coverage of numerous variables that were considered; the use of a more specific and eventually more valid measure of occupational outcome; and the use of recursive partitioning.

Classically in prospective cohort studies, subjects must be free of the disease at the beginning of observation, so that only new (incident) cases are observed and the directionality of associations is non-equivocal. Since only about 1 in 5 back-pain patients consulting in primary care settings have never had back pain before, a so-called inception cohort study of incident cases would require great resources and include only subjects with homogeneous characteristics, which would limit its external validity. Primary care physicians meet with a heterogeneous population of back-pain patients (most with recurrent or persistent pain) for whom they must give a prognosis. A prediction tool that works for a subgroup of patients only is unlikely to be useful to clinicians, especially if the subgroup includes only a minority of their clients.

In this study, several measures of the type and severity of pain were used as potential predictors. If these variables had been important to the prognosis, they would have been retained. Our model is thus applicable to all sorts of back-pain problems, as seen in day-to-day primary care practice. It can be used systematically to assist the physician in deciding the best allocation of clinical resources for



**Fig. 2: Progress among study participants toward return to work in good health.** Note that subjects could not go back into the “failure” group; this category could thus only diminish over time.



**Fig. 3: Clinical algorithm to predict an outcome at 2 years of return to work in good health (RWGH) among workers consulting in primary care settings for back pain.** All values shown are percentages. High-probability categories in each group, as were used to calculate the measures of validity, are circled. Note that the "failure" category includes lack of successful return to work at 2 years of follow-up, either with no attempt to return or despite 1 or more attempts to return to work. CI = confidence interval, Q = question. Q1 is item 15 of the Fear Avoidance Beliefs Questionnaire. Questions 5-7 are from the Roland-Morris Disability Questionnaire: Q5 is item 2; Q6, item 22; and Q7, item 18.

Example 1. Mr. Jones answered "Yes" to question 1, "No" to question 3, "7" to question 4 and "No" to question 5. (Notice that because Mr. Jones said "Yes" to question 1, question 2 is not useful.) His estimated probability of success at 2 years in returning to work in good health is 84%. The clinician will reassure him and use a conservative approach. A rapid return to normal activities is the objective.

Example 2. Mr. Smith answered "No" to question 1 and "Yes" to question 2. (Questions 3-7 are unnecessary for Mr. Smith.) He thus appears to have a particularly high probability (46%) of failure to return to work in good health by 2 years. The clinician may wish to refer him to a specialized rehabilitation program.

Example 3. Mrs. Watson answered "Yes" to question 1, "No" to question 3, "8" to question 4, "Yes" to questions 5 and 6, and "No" to question 7. (Again, question 2 is not needed in Mrs. Watson's case.) Her probability of either success or partial success in returning to work in good health by 2 years is quite high (50% + 45% = 95%). The clinician could ask to see her again and eventually refer her to occupational health services to monitor and improve her work conditions. Keeping the patient at work is the objective.

patients with back pain (see the example cases accompanying Fig. 3).

Reproducibility of predictors is an important consideration in building a prognostic instrument.<sup>68</sup> The fact that 5 of the variables included in our final model are items drawn from well-validated measurement instruments is reassuring. The 2 other items, radiating pain and previous back surgery, are relatively “hard” events, measurement of which is likely to be highly reproducible.

Because the baseline interviews were conducted some 3 weeks after the related medical consultation, it is possible that some variables may have changed during that period,

Predicted 2-year outcome	Actual 2-year outcome		Totals
	Failure or failure after attempt	Success or partial success	
Failure or failure after attempt	a (45)	b (100)	a + b (145)
Success or partial success	c (12)	d (180)	c + d (192)
Totals	a + c (57)	b + d (280)	(337*)

Sensitivity:  $a \div (a + c) = 45/57 = 79.0\%$   
 Specificity:  $d \div (b + d) = 180/280 = 64.3\%$   
 Positive predictive value:  $a \div (a + b) = 45/145 = 31.0\%$   
 Negative predictive value:  $d \div (c + d) = 180/192 = 93.8\%$   
 Prevalence of adverse outcome:  $(a + c) \div (a + b + c + d) = 57/337* = 16.9\%$

\*17 subjects had missing values for 1 or more predictors.

**Fig. 4: Example of calculations of the measures of validity presented in Table 3.**

improving the predictive potential over data that would have been collected, had the interviews been held at the time of consultation. However, variables measured at 6 weeks and 12 weeks offered no better prediction than the baseline measures, which is reassuring.

Traditionally, physicians look for clinical decision rules that have high positive predictive value, and attribute generally less importance to negative predictive value. However, it is the nature of the outcome that must determine the most important measures of validity for a given clinical decision tool. Considering the frequency of back pain and the resources that are spent on benign cases, an instrument that allows identification of a group of subjects who are at low risk of adverse outcomes may be quite useful.

This article has been peer reviewed.

From Unité de recherche en santé des populations, Centre de recherche du Centre hospitalier affilié universitaire de Québec (Dionne, Larocque); the Local Community Health Care Center (CLSC) Haute-Ville-des-Rivières, Quebec City, Que. (Bourbonnais); Département de réadaptation, Faculté de médecine, Université Laval, Sainte-Foy, Que. (Dionne, Bourbonnais, Frémont); Département d'épidémiologie, de biostatistique et de santé au travail, Université McGill, and Régie régionale de la santé et des services sociaux de Montréal-centre, Montréal, Que. (Rossignol, Stock); and Institut national de santé publique du Québec, Montréal, Que. (Stock).

Competing interests: None declared.

Contributors: Clermont Dionne initiated the study. Clermont Dionne, René Bourbonnais, Pierre Frémont, Michel Rossignol and Susan Stock designed the study protocol. All of the authors discussed core ideas, participated in analysis and interpretation of the data, and contributed to the writing of the paper under Clermont Dionne's lead. Clermont Dionne and Isabelle Larocque coordinated the study and data collection. All of the authors are guarantors for the paper.

Acknowledgements: We thank all of the study participants, the staff of the Unité de recherche en santé des populations and all of the research assistants who worked on this study. Thanks also to physicians Stéphane Bergeron, Alexandra Dansereau, Georges Dufresne, Louis Larue, Natalie Le Sage, Jean Maziade and Jean Ouellet for their help with the recruitment of subjects. Special thanks to Arie Nouwen for his help with the questionnaire on self-efficacy, Julie Soucy for her participation in the coordination of the study and Eric Demers for his contribution to the statistical analyses.

This study was supported by grant 97-061 from the Quebec Institute for Occupational Safety and Health (Institut de recherche Robert-Sauvé en santé et en sécurité du travail du Québec), which did not interfere in any way in the scientific and publication processes. Clermont Dionne and René Bourbonnais are Quebec Health Research Fund (Fonds de la recherche en santé du Québec) Scholars.

**Table 3: Measures of validity\* for the final predictive model of the 2-year outcome “return to work in good health” in the 2 samples of study participants**

Validity measure	% of training sample, n = 354			% of validation sample, n = 506		
	Failure or failure after attempt	Partial success	Partial success + failure or failure after attempt	Failure or failure after attempt	Partial success	Partial success + failure or failure after attempt
Classification error <sup>†</sup>		37.0			40.5	
Sensitivity	79	64	73	74	60	69
Specificity	64	60	71	62	59	63
Positive predictive value	31	32	66	33	29	57
Negative predictive value	94	85	78	91	84	74

\*Calculated from 2x2 tables comparing each definition of outcome in the heading (the “diseased” group) to the other remaining category or categories (the “non-diseased” group, which always includes at least the “success” category). An example of a set of actual calculations is provided in Fig. 4.

<sup>†</sup>Although useful for comparison purposes, classification error rates tend to be biased toward the prevalence of the outcome and should therefore be interpreted with caution.

## References

- Hart LG, Deyo RA, Cherkin DC. Physician office visits for low back pain: frequency, clinical evaluation, and treatment patterns from a US national survey. *Spine* 1995;20:11-9.
- Cypress BK. Characteristics of physician visits for back symptoms: a national perspective. *Am J Public Health* 1983;73(4):389-95.
- Rapaport J, Jacobs P, Bell NR, Klarenbach S. Refining the measurement of the economic burden of chronic diseases in Canada. *Chronic Dis Can Winter* 2004;25(1):13-21.
- Druss B. The most expensive medical conditions in America. *Health Affairs* 2002;21(4):105-11.
- Bigos S, Bowyer O, Braen G, et al. *Acute low back problems in adults*. [Clinical Practice Guideline no 14.] Rockville (MD): Agency for Health Care Policy and Research, Public Health Service, US Department of Health and Human Services; 1994. Publication no 95-0642.
- Deyo RA, Weinstein JN. Low back pain. *N Engl J Med* 2001;344(5):363-70.
- Deyo RA, Cherkin D, Conrad D, Volinn E. Cost, controversy, crisis: low back pain and the health of the public. *Annu Rev Public Health* 1991;12:141-56.
- Volinn E, Van Koevering D, Loeser JD. Back sprain in industry: the role of socioeconomic factors in chronicity. *Spine* 1991;16(5):542-8.
- Gatchel RJ, Polatin PB, Mayer TG. The dominant role of psychosocial risk factors in the development of chronic low back pain disability. *Spine* 1995;20(24):2702-9.
- Spitzer WO. Scientific approach to the assessment and management of activity-related spinal disorders: a monograph for clinicians. Report of the Quebec Task Force on Spinal Disorders. *Spine* 1987;12(7 Suppl):S1-59.
- Abenhaim L, Rossignol M, Gobeille D, Bonvalot Y, Fines P, Scott S. The prognostic consequences in the making of the initial medical diagnosis of work-related back injuries. *Spine* 1995;20(7):791-5.
- De Girolamo G. Epidemiology and social costs of low back pain and fibromyalgia. *Clin J Pain* 1991;7(Suppl 1):S1-7.
- Spengler DM, Bigos SJ, Martin NA, Zeh J, Fisher L, Nachemson A. Back injuries in industry: a retrospective study. I. Overview and cost analysis. *Spine* 1986;11(3):241-5.
- Hashemi L, Webster BS, Clancy EA, Volinn E. Length of disability and cost of workers' compensation low back pain claims. *J Occup Environ Med* 1997;39(10):937-45.
- Butterfield PG, Spencer PS, Redmond N, Feldstein A, Perrin N. Low back pain: predictors of absenteeism, residual symptoms, functional impairment, and medical costs in Oregon workers' compensation recipients. *Am J Ind Med* 1998;34(6):559-67.
- Van der Giezen AM, Bouter LM, Nijhuis FJ. Prediction of return-to-work of low back pain patients sicklisted for 3-4 months. *Pain* 2000;87(3):285-94.
- Feuerstein M, Berkowitz SM, Huang GD. Predictors of occupational low back disability: implications for secondary prevention. *J Occup Environ Med* 1999;41(12):1024-31.
- Boos N, Semmer N, Elfering A, Schade V, Gal I, Zanetti M, et al. Natural history of individuals with asymptomatic disc abnormalities in magnetic resonance imaging: predictors of low back pain-related medical consultation and work incapacity. *Spine* 2000;25(12):1484-92.
- Viihari-Juntura E, Takala E, Riihimäki H, Martikainen R, Jappinen P. Predictive validity of symptoms and signs in the neck and shoulders. *J Clin Epidemiol* 2000;53(8):800-8.
- Linton SJ, Hallden K. Can we screen for problematic back pain? A screening questionnaire for predicting outcome in acute and subacute back pain. *Clin J Pain* 1998;14(3):209-15.
- Lancourt J, Kettelhut M. Predicting return to work for lower back pain patients receiving worker's compensation. *Spine* 1992;17(6):629-40.
- Cats-Baril WL, Frymoyer JW. Identifying patients at risk of becoming disabled because of low-back pain: the Vermont Rehabilitation Engineering Center predictive model. *Spine* 1991;16(6):605-7.
- Lehmann TR, Spratt KF, Lehmann KK. Predicting long-term disability in low back injured workers presenting to a spine consultant. *Spine* 1993;18:1103-12.
- Hogg-Johnson S, Cole DC. Early prognostic factors for duration on temporary total benefits in the first year among workers with compensated occupational soft tissue injuries. *Occup Environ Med* 2003;60(4):244-53.
- McIntosh G, Frank J, Hogg-Johnson S, Bombardier C, Hall H. Prognostic factors for time receiving workers' compensation benefits in a cohort of patients with low back pain. *Spine* 2000;25(2):147-57.
- Baldwin ML, Johnson WG, Butler RJ. The error of using returns-to-work to measure the outcomes of health care. *Am J Ind Med* 1996;29:632-41.
- Atlas SJ, Deyo RA, Patrick DL, Convery K, Keller RB, Singer DE. The Quebec Task Force classification for spinal disorders and the severity, treatment, and outcomes of sciatica and lumbar spinal stenosis. *Spine* 1996;21:2885-92.
- Roland M, Morris R. A study of the natural history of back pain. Part I: development of a reliable and sensitive measure of disability in low-back pain. *Spine* 1983;8(2):141-4.
- Waddell G. *The back pain revolution*. Edinburgh: Churchill Livingstone; 1998. p. 39-40.
- Jensen MP, Strom SE, Turner JA, Romano JM. Validity of the Sickness Impact Profile Roland Scale as a measure of dysfunction in chronic pain patients. *Pain* 1992;50(2):157-62.
- Beurskens AJ, de Vet HC, Koke AJ, van der Heijden GJ, Knipschild PG. Measuring the functional status of patients with low back pain: assessment of the quality of four disease-specific questionnaires. *Spine* 1995;20(9):1017-28.
- Roland M, Fairbank J. The Roland-Morris Disability Questionnaire and the Oswestry Disability Questionnaire. *Spine* 2000;25(24):3115-24.
- Riddle DL, Stratford PW, Binkley JM. Sensitivity to change of the Roland-Morris Back Pain Questionnaire: part 2. *Phys Ther* 1998;78(11):1197-207.
- Bolton JE. Accuracy of recall of usual pain intensity in back pain patients. *Pain* 1999;83(3):533-9.
- Strong J, Ashton R, Chant D. Pain intensity measurement in chronic low back pain. *Clin J Pain* 1991;7(3):209-18.
- Waddell G, Newton M, Henderson I, Somerville D, Main C. A fear-avoidance beliefs questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain* 1993;52:157-68.
- Jacob T, Baras M, Zeev A, Epstein L. Low back pain: reliability of a set of pain measurement tools. *Arch Phys Med Rehabil* 2001;82(6):735-42.
- Derogatis LR. *Symptoms Checklist-90. Administration, scoring and procedures manual for the revised version*. Baltimore: Clinical Psychometric Research; 1977.
- Dionne CE, Bourbonnais R, Frémont P, et al. Predictors of return to work in good health among back pain patients in primary care settings: 3-month outcome in the RAMS-Prognosis Study [abstract]. International Forum V on Primary Care Research on Low Back Pain, 2002; Montreal.
- Dozois DJA, Dobson KS, Wong M, Hughes D, Long A. Factors associated with rehabilitation outcome in patients with low back pain (LBP): prediction of employment outcome at 9-month follow-up. *Rehabil Psychol* 1995;40(4):243-59.
- Severeijns R, Vlaeyen JW, van den Hout MA, Weber WE. Pain catastrophizing predicts pain intensity, disability, and psychological distress independent of the level of physical impairment. *Clin J Pain* 2001;17(2):165-72.
- Nickel R, Egle UT, Eysel P, Rompe JD, Zollner J, Hoffmann SO. Health-related quality of life and somatization in patients with long-term low back pain: a prospective study with 109 patients. *Spine* 2001;26(20):2271-7.
- Shutty MS Jr, DeGood DE, Schwartz DP. Psychological dimensions of distress in chronic pain patients: a factor analytic study of Symptom Checklist-90 responses. *J Consult Clin Psychol* 1986;54(6):836-42.
- Karasek RA. *Job Content Questionnaire and User's Guide*. Los Angeles (CA): Department of Industrial and System Engineering, University of Southern California; 1985.
- Brisson C, Blanchette C, Guimont C, Dionne G, Moisan J, Vézina M. Reliability and validity of the French version of the 18-item Karasek Job Content Questionnaire. *Work Stress* 1998;12(4):322-36.
- Bigos SJ, Battie MC, Spengler DM, Fisher LD, Fordyce WE, Hansson TH, et al. A prospective study of work perceptions and psychosocial factors affecting the report of back injury [published erratum appears in *Spine* 1991;16(6):688]. *Spine* 1991;16(1):1-6.
- Friedrich M, Cermak T, Heiller I. Spinal troubles in sewage workers: epidemiological data and work disability due to low back pain. *Int Arch Occup Environ Health* 2000;73(4):245-54.
- Reigo T, Tropp H, Timpka T. Absence of back disorders in adults and work-related predictive factors in a 5-year perspective. *Eur Spine J* 2001;10(3):215-20; discussion 221.
- Williams DA, Feuerstein M, Durbin D, Pezzullo J. Health care and indemnity costs across the natural history of disability in occupational low back pain. *Spine* 1998;23(21):2329-36.
- Williams RA, Pruitt SD, Doctor JN, Epping-Jordan JE, Wahlgren DR, Grant I, et al. The contribution of job satisfaction to the transition from acute to chronic low back pain. *Arch Phys Med Rehabil* 1998;79(4):366-74.
- Loisel P, Durand P, Gosselin L, Simard R, Turcotte J, editors. *La clinique des maux de dos — un modèle de prise en charge en prévention de la chronicité*. Québec: Institut de recherche en santé et en sécurité du travail du Québec; mai 1996.
- Siegrist J, Starke D, Chandola T, Godin I, Marmot M, Niedhammer I, et al. The measurement of effort-reward imbalance at work: European comparisons. *Soc Sci Med* 2004;58(8):1483-99.
- Niedhammer I, Siegrist J, Landre MF, Goldberg M, Leclerc A. [Psychometric properties of the French version of the Effort-Reward Imbalance model]. *Rev Epidemiol Sante Publique* 2000;48(5):419-37.
- Cherkin D, Deyo RA, Berg AO, Bergman JJ, Lishner DM. Evaluation of a physician education intervention to improve primary care for low-back pain. I. Impact on physicians. *Spine* 1991;16(10):1168-72.
- Von Korff M, Barlow W, Cherkin D, Deyo RA. Effects of practice style in managing back pain. *Ann Intern Med* 1994;121(3):187-95.
- Rosenstiel AK, Keefe FJ. The use of coping strategies in chronic low back pain patients: relationship to patient characteristics and current adjustment. *Pain* 1983;17:33-44.
- Cedraschi C, Nordin M, Nachemson AL, Vischer TL. Health care providers should use a common language in relation to low back pain patients. *Baillieres Clin Rheumatol* 1998;12:1-15.
- Bandura A. Self-efficacy: toward a unifying theory of behavioral change. *Psychol Rev* 1977;84(2):191-215.
- Wallston KA, Wallston BS, DeVellis R. Development of the Multidimensional

- Health Locus of Control (MHLC) Scales. *Health Educ Monogr* 1978;6(2):160-70.
60. Talbot F, Nouwen A, Gauthier J. Is health locus of control a 3-factor or a 2-factor construct? *J Clin Psychol* 1996;52(5):559-68.
61. Vallerand RJ. Vers une méthode de validation trans-culturelle des questionnaires psychologiques: implications pour la recherche en langue française. *Can Psychol* 1989;30:662.
62. Kleinbaum DG, Kupper LL, Muller KE. *Applied regression analysis and other multivariable methods*. 2nd ed. Boston: PWS-Kent Publishing Company; 1988. p. 386-482.
63. Breiman L, Friedman JH, Olshen RA, Stone CJ. *Classification and regression trees*. Pacific Grove (CA): Wadsworth & Brooks; 1984. 368 p.
64. Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology* 1982;143:29-36.
65. Dionne CE, Koepsell TD, Von Korff M, Deyo RA, Barlow WE, Checkoway H. Predicting long-term functional limitations among back pain patients in primary care settings. *J Clin Epidemiol* 1997;50:31-43.
66. McIntosh G, Frank J, Hogg-Johnson S, Hall H, Bombardier C. Low back pain prognosis: structured review of the literature. *J Occup Rehabil* 2000;10(2):101-15.
67. Thomas E, Silman AJ, Croft PR, Papageorgiou AC, Jayson MI, Macfarlane GJ. Predicting who develops chronic low back pain in primary care: a prospective study. *BMJ* 1999;318(7199):1662-7.
68. Laupacis A, Sekar N, Stiell IG. Clinical prediction rules. A review and suggested modifications of methodological standards. *JAMA* 1997;277(6):488-94.

**Correspondence to:** Dr. Clermont E. Dionne, Unité de recherche en santé des populations, Centre de recherche du Centre hospitalier affilié universitaire de Québec, Hôpital du Saint-Sacrement, 1050, chemin Ste-Foy, Québec QC G1S 4L8; fax 418 682-7949; clermont.dionne@uresp.ulaval.ca

## Appendix 1: List of constructs measured and considered as potential predictors

### Sociodemographic factors

- Age
- Current financial problems because of back pain
- Education
- Ethnic origin
- Gender
- Gross annual family income
- Living with spouse
- Maternal language
- Number of household members
- Number of children under 5 years of age

### Anthropometric factors

- Body mass index (kg/m<sup>2</sup>)
- Hand dominance
- Height
- Weight

### Psychological factors

- Aggressiveness
- Exposure to stressful events during the past 12 months: changed job for worse, divorced or separated, financial problems, got gravely ill, significant other became gravely ill, death of spouse, death of other significant person)
- Fear-avoidance beliefs — activity
- Fear-avoidance beliefs — work
- Health locus of control (internal, external, chance)
- Pain-control strategies
- Psychological trauma before age 18
- Self-efficacy with regard to work capacity
- Symptoms of depression
- Symptoms of somatization
- Social support outside work

### Health behaviour factors

- Alcoholism
- Cigarette smoking
- Coffee and tea consumption
- Physical activity and sports participation when not in pain
- Regular meditation or relaxation
- Sleep during past 2 weeks
- Usual duration of sleep

### Clinical factors, history of back pain

- Age at menopause
- Average pain over the past 6 months
- Current functional limitations
- Current pain level
- Diagnostic category
- Duration of current episode at time of consultation
- Duration of pain, past 6 months
- Number of previous surgeries for back pain
- Number of days with pain since index consultation
- Number of times gave birth
- Pain radiating to upper or lower limb?
- Previous back pain episode(s) affecting work capacity and, if any, when?
- Reduction of daily life activities because of back pain, over the past 6 months
- Reduction of work capacity because of back pain, past 6 mo
- Self-reported health status
- Self-reported type of back pain
- Site of pain
- Spitzer classification<sup>10,70</sup>
- Worst pain (level) in past 6 months
- Number of years since onset of back pain

**Appendix 1: List of constructs measured and considered as potential predictors** (continued)**Dramatization of diagnosis**

- “The doctor asked you to get an x-ray, a CT scan, myelography, MRI (magnetic resonance imaging), or other radiological examination”
- “The doctor told you to stay in bed” (and no. of days recommended)
- “The doctor asked you to schedule a check-back visit”
- “The doctor mentioned that you might need surgery for your back problem”

**Utilization of health services**

- Duration of index medical consultation
- Number of visits to the same clinic during the past 12 months
- Number of hospitalizations for back pain during the past 12 months
- Perception that doctor’s diagnosis of back problem is correct
- Satisfaction with health services since the index consultation
- “The doctor asked you to undress for the physical examination of your back”
- “The doctor, physical therapist, or another health professional helped you learn exercises for your back to help in your recovery”
- “The doctor showed interest in you as a person”
- “The doctor asked about your lifestyle and stress that may affect your back pain”
- “The doctor discussed options with you, asked your opinion, and offered choices about the care of your back pain”
- “The doctor relieved any concern about the cause of your back pain”
- “Your visit to the doctor helped you feel that you could control your back pain”
- “You were treated like your back pain was an important problem, and not ‘all in your head’”
- “The doctor listened carefully while you described your back problem”
- “The doctor took time to answer all your questions about your back pain”
- “The doctor asked you about your occupation and took time to give you advice”
- “Your doctor, physical therapist or another professional showed you how to work to protect your back from injury”
- Weight of the medical file

**Occupational factors**

- Adaptation of work environment is possible
- Change of job or employer in the past 12 months
- Company size
- Current financial compensation for back pain
- Days of sick-leave permitted without a medical certificate
- Declared occupational disorder
- Effort–reward imbalance (Siegrist’s model)
- Exposure to cold at work
- Job decision latitude (Karasek’s model)
- Job satisfaction
- Job seniority
- Job status at time of consultation: permanent, occasional/seasonal, self-employed, other
- Likelihood of losing job during the next 2 years
- Number of days off work or with modified duties or schedule, over the past 12 months
- Number of jobs held
- Number of previous periods of sick-leave, modified duties, or modified schedule because of back pain over the past 10 years
- Pace of work
- Physical demands of job: sitting and standing, forward and lateral trunk flexion, rotation of the trunk, manipulation of charges, efforts made with tools, lifting or transportation of persons, whole-body vibration, physical strength
- Previous financial compensation for back pain
- Psychological demands of work (Karasek’s model)
- Responsibility for household chores
- Satisfaction concerning possibilities of work adaptation
- Social support at work
- “Supervisor takes back pain seriously”
- Time to commute to work
- Unionized job
- “When did you modify work or stop working for the current episode of back pain?”
- Work schedule: full-time, part-time
- Work shift: day, evening, night, variable
- Work status at time of consultation: at regular job, completely off work, modified duties, modified schedule, modified job assignment