

CLINICAL IMAGES

Syphilitic chancre of the mouth

Jianjun Qiao MD PhD, Hong Fang MD MPhil



Figure 1: A 1-cm nontender, nonpurulent, oval ulcer with clean base and raised rolled border on the lip of a 24-year-old man.

A 24-year-old man presented with a three-week history of a 1-cm painless, ulcerative lesion on his lower lip (Figure 1) and a three-day history of symmetrically distributed nonpruritic macules on his trunk (Figure 2) and limbs. He was HIV-negative and was otherwise healthy. He had performed unprotected orogenital sex with a female partner about three weeks before the onset of the ulcer. On examination, generalized nontender lymphadenopathy was noted along with the rash and chancre. A serologic test for syphilis showed a reactive rapid plasma reagin test result (titer 1:64) and a positive agglutination test result for *Treponema pallidum*. Syphilis was diagnosed, with findings consistent with both primary (oral chancre) and secondary (lymphadenopathy and macular rash) disease. He received treatment with intramuscular injection of penicillin G benzathine. The mucocutaneous lesions improved rapidly and resolved at the end of the treatment.

Transmission of acquired syphilis occurs mostly through sexual intercourse.^{1,2} Syphilitic chancre develops 3–90 days after exposure (mean 3 weeks),^{1,2} with genitalia as the most common inoculation sites.² Extragenital chancres occur in 12%–14% of patients with primary syphilis, the oral mucosa being the most frequent location as a consequence of orogenital con-



Figure 2: Violaceous macules on the patient's trunk.

tact.^{1,3} Diagnosis of syphilitic chancre is based on the patient's history of orogenital sexual contact, a reasonable incubation period, clinical features and results of serologic tests for syphilis. The differential diagnosis includes chancroid, herpes simplex, tuberculous chancre, deep mycoses, squamous cell carcinoma, traumatic ulcer, aphthous stomatitis and Behçet syndrome.³

Without treatment, syphilitic chancre spontaneously resolves in two to eight weeks. The secondary stage develops 2 to 12 weeks (mean 8 weeks) after inoculation.² The secondary stage may overlap with the primary stage, as seen in our patient.

References

1. Ficarra G, Carlos R. Syphilis: the renaissance of an old disease with oral implications. *Head Neck Pathol* 2009;3:195-206.
2. Dylewski J, Duong M. The rash of secondary syphilis. *CMAJ* 2007;176:33-5.
3. Alam F, Argiriadou AS, Hodgson TA, et al. Primary syphilis remains a cause of oral ulceration. *Br Dent J* 2000;189:352-4.

Competing interests: None declared.

This article has been peer reviewed.

Affiliation: From the Department of Dermatology, The First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou, People's Republic of China

Correspondence to: Dr. Hong Fang, fanghongzy@sina.com

CMAJ 2011. DOI:10.1503/cmaj.110664