

## Latency period of radiation-induced cancer

Some information in this article is flawed, as the authors have not properly allowed for the latency period for cancer induction by radiation.<sup>1</sup> A period of one year after the incident radiation may be insufficient. The latency period for induction of leukemia is 5–7 years, and for solid tumours is at least 10 years, so cancers occurring earlier than this should be considered to be naturally occurring rather than induced by radiation exposure.<sup>2</sup>

**Donald K. Fraser MD**

Associated Radiologists of Saskatoon,  
Saskatoon, Sask.

### Reference

1. Eisenberg MJ, Afilalo J, Lawler PR, et al. Cancer risk related to low-dose ionizing radiation from cardiac imaging in patients after acute myocardial infarction. *CMAJ* 2011;183:430-6.
2. Hall EJ. *Radiobiology for the radiologist*. 5th ed. Philadelphia (PA): Lippincott Williams & Wilkins; 2000:p. 149.

*CMAJ* 2011. DOI:10.1503/cmaj.111-2088

### The authors respond:

With respect to the shorter latency period than was observed in the Life Span Study (LSS),<sup>1</sup> we agree that this is an area that needs to be explored further. Several possibilities could explain this finding. The patient population in our cohort had about a twofold higher incidence of cancer — independent of exposure to low-dose ionizing radiation<sup>2</sup> — than nationally quoted statistics on cancer incidence rates.<sup>3</sup> The patients in our cohort were more at risk for cancer (perhaps because of smoking or obesity), and as such the latency time to cancer might have been shortened. To control for this, we used a time-dependent regression analysis, which incorporated a time-lag covariant to allow for cancer “induction” time following exposure. When sensitivity analyses were performed looking at time lags up to five years, there was no appreciable effect on the results. Nonetheless, we agree that important questions about the observed time lag in our study need

to be explained further, and we are attempting to define this more extensively in on-going studies.

**Mark J. Eisenberg MD MPH**

**Louise Pilote MD MPH PhD**

**Patrick R. Lawler MD**

**Jonathan Afilalo MD MSc**

McGill University, Montréal, Que.

### References

1. National Research Council of the National Academies, Committee to Assess Health Risks from Exposure to Low Levels of Ionizing Radiation. *Biological effects of ionizing radiation — BEIR VII phase 2*. Washington (DC): National Academies Press; 2006.
2. Eisenberg MJ, Afilalo J, Lawler PR, et al. Cancer risk related to low-dose ionizing radiation from cardiac imaging in patients after acute myocardial infarction. *CMAJ* 2011;183:430-6.
3. Canadian Cancer Society's Steering Committee. *Canadian cancer statistics 2009*. Toronto (ON): Canadian Cancer Society; 2009. p. 24.

*CMAJ* 2011. DOI:10.1503/cmaj.111-2089

## Don't select medical students — convince them

Scott and colleagues presented a study of high quality, addressing an important topic.<sup>1</sup> The authors encourage medical faculties to increase their output of family medicine residents by selecting and targeting students based on sociodemographic and attitudinal variables of predictive value. Such an approach would require the acquisition and storage of personal information. To our mind, this is concerning, given ethical consideration and data privacy. The education level of parents, relationship status and societal attitudes should not influence admission to medical school or course and content of medical education. We see a bigger benefit in understanding the variables that influence the decision-making process in the course of studies. Recent publications show that positive family medicine experiences, longitudinally implemented in the curriculum, positively influence the attractiveness of family medicine as a career.<sup>2-5</sup> Obligatory and longer family medicine clerkships and exposure to role models influence students committed to primary care as well as those who are undecided.<sup>3</sup> Sev-

eral studies describe the positive effect of rural training experiences.<sup>4,6</sup> So don't select students — convince them.

**Tobias Deutsch MSc**

**Petra Hönigschmid**

**Uta Wippermann**

**Thomas Frese MD**

**Hagen Sandholzer MD PhD**

Department of Primary Care, Leipzig  
Medical School, Leipzig, Germany

### References

1. Scott I, Gowans M, Wright B. Determinants of choosing a career in family medicine. *CMAJ* 2011; 83: E1-8.
2. Bunker J, Shadbolt N. Choosing general practice as a career — the influences of education and training. *Aust Fam Physician* 2009;38:341-4.
3. Bennett KL, Phillips JP. Finding, recruiting, and sustaining the future primary care physician workforce: a new theoretical model of specialty choice process. *Acad Med* 2010;85(Suppl):S81-8.
4. Dick JF 3rd, Wilper AP, Smith S, et al. The effect of rural training experiences during residency on the selection of primary care careers: a retrospective cohort study from a single large internal medicine residency program. *Teach Learn Med* 2011;23 (1):53-7.
5. Shadbolt N, Bunker J. Choosing general practice — a review of career choice determinants. *Aust Fam Physician* 2009;38:53-5.
6. Barrett FA, Lipsky MS, Lutfiyya MN. The impact of rural training experiences on medical students: a critical review. *Acad Med* 2011;86:259-63.

*CMAJ* 2011. DOI:10.1503/cmaj.111-2084

## Electroconvulsive therapy article unbalanced

*CMAJ* represents the views of and informs all Canadian physicians. Thus psychiatrists were dismayed by the sensationalistic article about the Food and Drug Administration's review of the classification of electroconvulsive therapy (ECT) devices.<sup>1</sup>

The stigmatizing photo accompanying the article offers a frightening and unrealistic portrayal of the administration of ECT. Stigma affecting patients who have mental health issues is far reaching<sup>2</sup> and *CMAJ* should show a high level of sensitivity to this issue.

The article observes that most recipients of ECT are women but neglects to give context to this comment. This allows the inference of impropriety whereas major depressive disorder, the primary condition treated with ECT, is highly prevalent among women.