Latency period of radiation-induced cancer

Some information in this article is flawed, as the authors have not properly allowed for the latency period for cancer induction by radiation.¹ A period of one year after the incident radiation may be insufficient. The latency period for induction of leukemia is 5–7 years, and for solid tumours is at least 10 years, so cancers occurring earlier than this should be considered to be naturally occurring rather than induced by radiation exposure.²

Donald K. Fraser MD

Associated Radiologists of Saskatoon,
Saskatoon, Sask.

Reference

1. Eisenberg MJ, Afilalo J, Lawler PR, et al. Cancer risk related to low-dose ionizing radiation from cardiac imaging in patients after acute myocardial infarction. *CMAJ* 2011;183:430-6.
2. Hall EJ. *Radiobiology for the radiologist*. 5th ed. Philadelphia (PA): Lippincott Williams & Wilkins; 2000:p. 149.

CMAJ 2011. DOI:10.1503/cmaj.111-2088

The authors respond:

With respect to the shorter latency period than was observed in the Life Span Study (LSS),¹ we agree that this is an area that needs to be explored further. Several possibilities could explain this finding. The patient population in our cohort had about a twofold higher incidence of cancer — independent of exposure to low-dose ionizing radiation² — than nationally quoted statistics on cancer incidence rates.³ The patients in our cohort were more at risk for cancer (perhaps because of smoking or obesity), and as such the latency time to cancer might have been shortened. To control for this, we used a time-dependent regression analysis, which incorporated a time-lag covariant to allow for cancer “induction” time following exposure. When sensitivity analyses were performed looking at time lags up to five years, there was no appreciable effect on the results. Nonetheless, we agree that important questions about the observed time lag in our study need

to be explained further, and we are attempting to define this more extensively in on-going studies.

Mark J. Eisenberg MD MPH
Louise Pilote MD MPH PhD
Patrick R. Lawler MD
Jonathan Afilalo MD MSc
McGill University, Montréal, Que.

References

1. National Research Council of the National Academies, Committee to Assess Health Risks from Exposure to Low Levels of Ionizing Radiation. *Biological effects of ionizing radiation — BEIR VII phase 2*. Washington (DC): National Academies Press; 2006.
2. Eisenberg MJ, Afilalo J, Lawler PR, et al. Cancer risk related to low-dose ionizing radiation from cardiac imaging in patients after acute myocardial infarction. *CMAJ* 2011;183:430-6.
3. Canadian Cancer Society's Steering Committee. *Canadian cancer statistics 2009*. Toronto (ON): Canadian Cancer Society; 2009. p. 24.

CMAJ 2011. DOI:10.1503/cmaj.111-2089

Don't select medical students — convince them

Scott and colleagues presented a study of high quality, addressing an important topic.¹ The authors encourage medical faculties to increase their output of family medicine residents by selecting and targeting students based on sociodemographic and attitudinal variables of predictive value. Such an approach would require the acquisition and storage of personal information. To our mind, this is concerning, given ethical consideration and data privacy. The education level of parents, relationship status and societal attitudes should not influence admission to medical school or course and content of medical education. We see a bigger benefit in understanding the variables that influence the decision-making process in the course of studies. Recent publications show that positive family medicine experiences, longitudinally implemented in the curriculum, positively influence the attractiveness of family medicine as a career.²⁻⁵ Obligatory and longer family medicine clerkships and exposure to role models influence students committed to primary care as well as those who are undecided.³ Sev-

eral studies describe the positive effect of rural training experiences.^{4,6} So don't select students — convince them.

Tobias Deutsch MSc
Petra Hönigsmid
Uta Wippermann
Thomas Frese MD
Hagen Sandholzer MD PhD
Department of Primary Care, Leipzig
Medical School, Leipzig, Germany

References

1. Scott I, Gowans M, Wright B. Determinants of choosing a career in family medicine. *CMAJ* 2011; 183: E1-8.
2. Bunker J, Shadbolt N. Choosing general practice as a career — the influences of education and training. *Aust Fam Physician* 2009;38:341-4.
3. Bennett KL, Phillips JP. Finding, recruiting, and sustaining the future primary care physician workforce: a new theoretical model of specialty choice process. *Acad Med* 2010;85(Suppl):S81-8.
4. Dick JF 3rd, Wilper AP, Smith S, et al. The effect of rural training experiences during residency on the selection of primary care careers: a retrospective cohort study from a single large internal medicine residency program. *Teach Learn Med* 2011;23 (1):53-7.
5. Shadbolt N, Bunker J. Choosing general practice — a review of career choice determinants. *Aust Fam Physician* 2009;38:53-5.
6. Barrett FA, Lipsky MS, Lutfiyya MN. The impact of rural training experiences on medical students: a critical review. *Acad Med* 2011;86:259-63.

CMAJ 2011. DOI:10.1503/cmaj.111-2084

Electroconvulsive therapy article unbalanced

CMAJ represents the views of and informs all Canadian physicians. Thus psychiatrists were dismayed by the sensationalistic article about the Food and Drug Administration's review of the classification of electroconvulsive therapy (ECT) devices.¹

The stigmatizing photo accompanying the article offers a frightening and unrealistic portrayal of the administration of ECT. Stigma affecting patients who have mental health issues is far reaching² and *CMAJ* should show a high level of sensitivity to this issue.

The article observes that most recipients of ECT are women but neglects to give context to this comment. This allows the inference of impropriety whereas major depressive disorder, the primary condition treated with ECT, is highly prevalent among women.

Why did this article include testimonials that had no credibility apart from being against ECT? Does *CMAJ* include the opinion of ideologues when discussing other medical devices or treatments? ECT is a life-saving and highly effective procedure for which there are clear guidelines.³

The Canadian Psychiatric Association stands by its stated position that “ECT has well defined indications, demonstrated efficacy and safety, well characterized side-effects, and established standards for practice. The decision to use ECT in the treatment of a patient is a medical one, based on the psychiatrist’s assessment of the patient’s illness and an evaluation of the merits and risks of ECT, compared with alternative treatments. Similar to all medical procedures, the use of ECT requires informed consent from the patient or a substituted decision maker.”²³

Nizar Ladha MD

President
Canadian Psychiatric Association, Ottawa,
Ont.

References

1. Benac N. United States reviews safety of electroconvulsive therapy. *CMAJ* 2011;183:E269-70.
2. Atzema CL, Schull MJ, Tu JV, et al. The effect of a charted history of depression on emergency triage and outcomes inpatients with acute myocardial infarction. *CMAJ* 2011;183:663-9.
3. Enns M, Reiss J, Chan P. Electroconvulsive therapy [position paper]. *Can J Psychiatry* 2011;55:1-11.

CMAJ 2011. DOI:10.1503/cmaj.111-2085

Cover of *CMAJ*

As a parent and a grandfather, I was glad that the Oct. 4, 2011, issue with the grossly diseased brain on the cover arrived at my home after my granddaughter had left from a visit with us. It is a disturbing image that as a physician and hospitalist on a brain injury unit I can understand. However, I do not think it belongs on the cover and I object to its placement there. I feel that there is a sensationalist approach to the use of this image and that the image did not serve the thoughtful article to which it was drawing attention. Nor did the image serve to advertise the medical profession as a group of people who would be likely to read an article unless it was

associated with lurid imagery. Surely, we can capture the attention of our colleagues with less troubling images and can see that our presentation of ourselves is reflected in the styling of our Canadian medical journal.

John Clement MD

Parkwood Hospital
London, Ont.

CMAJ 2011. DOI:10.1503/cmaj.111-2091

Addressing the root cause

Reinier and colleagues deserve congratulations for further establishing the link between low socioeconomic status and cardiovascular-related deaths.¹ However, that they restricted themselves to proposing solutions such as targeted training for cardiopulmonary resuscitation and placement of automated external defibrillators is disappointing. Surely, addressing low socioeconomic status itself has a role to play in reducing mortality from sudden cardiac arrest. As physicians, we have unique perspectives to offer in public policy discussions around poverty. We should not feel restrained in doing so.

Reference

1. Reinier K, Thomas E, Andrusiek DL, et al.; the Resuscitation Outcomes Consortium Investigators. Socioeconomic status and incidence of sudden cardiac arrest *CMAJ* 2011;183:1705-12.

Danyaal Raza

Fellow in Global Health
University of Toronto
Toronto, Ont.

CMAJ 2011. DOI:10.1503/cmaj.111-2090

Letters to the editor

In submitting a letter, you automatically consent to have it appear online and/or in print. All letters accepted for print will be edited by *CMAJ* for space and style. Most references and multiple authors’ names, full affiliations and competing interests will appear online only. (The full version of any letter accepted for print will be posted at cmaj.ca.)