

Effectiveness of disease-management programs for improving diabetes care: a meta-analysis

Clément Pimouguet MSc, Mélanie Le Goff MSc, Rodolphe Thiébaud MD PhD, Jean François Dartigues MD PhD, Catherine Helmer MD PhD

ABSTRACT

Background: We conducted a meta-analysis of randomized controlled trials to assess the effectiveness of disease-management programs for improving glycemic control in adults with diabetes mellitus and to study which components of programs are associated with their effectiveness.

Methods: We searched several databases for studies published up to December 2009. We included randomized controlled trials involving adults with type 1 or 2 diabetes that evaluated the effect of disease-management programs on glycated hemoglobin (hemoglobin A_{1c}) concentrations. We performed a meta-regression analysis to determine the effective components of the programs.

Results: We included 41 randomized controlled trials in our review. Across these trials, disease-management programs resulted in a significant reduction in hemoglobin A_{1c} levels (pooled standardized mean difference between intervention and control groups -0.38 [95% confidence interval -0.47 to -0.29], which corresponds to an absolute mean difference of

0.51%). The finding was robust in the sensitivity analyses based on quality assessment. Programs in which the disease manager was able to start or modify treatment with or without prior approval from the primary care physician resulted in a greater improvement in hemoglobin A_{1c} levels (standardized mean difference -0.60 v. -0.28 in trials with no approval to do so; $p < 0.001$). Programs with a moderate or high frequency of contact reported a significant reduction in hemoglobin A_{1c} levels compared with usual care; nevertheless, only programs with a high frequency of contact led to a significantly greater reduction compared with low-frequency contact programs (standardized mean difference -0.56 v. -0.30 , $p = 0.03$).

Interpretation: Disease-management programs had a clinically moderate but significant impact on hemoglobin A_{1c} levels among adults with diabetes. Effective components of programs were a high frequency of patient contact and the ability for disease managers to adjust treatment with or without prior physician approval.

Competing interests: See end of article for competing interests.

This article has been peer reviewed.

Correspondence to: Clément Pimouguet, clement.pimouguet@ispe.u-bordeaux2.fr

CMAJ 2011. DOI:10.1503/cmaj.091786

Despite well-established recommendations for diabetes care,¹⁻³ quality of care still needs to be improved. Although many nonpharmacologic strategies (patient education, psychological intervention, dietary education, self-monitoring and telemedicine) have been developed, their effectiveness is still unclear.⁴⁻⁶ “Disease management” is a structured, multifaceted intervention that includes several of the above-mentioned components. In two recent meta-analyses, disease management was associated with an improvement in glycemic control, as assessed by a mean reduction in hemoglobin A_{1c} concentration of 0.52% and 0.81%.^{7,8} Disease management seems to be more effective than single strategies such as clinician education, patient education or promotion of self-management.⁷

Because disease-management programs are heterogeneous, the effective components need to be identified to improve program implementation.

Previous studies have evaluated the efficacy of some program components.^{7,8} Independent medication changes by the disease manager appear to be particularly effective.⁷ However, other important factors such as the intensity of the intervention have not been previously evaluated.

We conducted a meta-analysis of randomized controlled trials (RCTs) involving adults with type 1 or 2 diabetes mellitus that evaluated the effect of disease-management programs on hemoglobin A_{1c} levels. We determined the effective components of the programs, considering both the type of component and the intensity of the intervention.

Methods

Definition of disease management

There is no consensual definition of disease management. According to the Care Continuum

Alliance (formerly the Disease Management Association of America), disease management “supports the physician or practitioner/patient relationship and plan of care; emphasizes prevention of exacerbations and complications utilizing evidence-based practice guidelines and patient empowerment strategies; and evaluates clinical, humanistic, and economic outcomes on an on-going basis with the goal of improving overall health” (www.carecontinuum.org/dm_definition.asp). To identify relevant studies for our meta-analysis, we adopted an operational definition based on the above definition, literature review and expert opinion.

We defined disease management as ongoing and proactive follow-up of patients that includes at least two of the following five components: patient education (dietary and exercise counselling, self-monitoring, and knowledge of disease and medication); coaching (the disease manager encourages the patient to overcome psychological or social barriers that impede autonomy or improvement in medication compliance); treatment adjustment (the disease manager is able to start or modify treatment with or without prior approval from the primary care physician); monitoring (the disease manager gets medical data from the patient); and care coordination (the disease manager reminds the patient about upcoming appointments or important aspects of self-care and informs the primary care physician about complications, treatment adjustment or therapeutic recommendations).

Literature search

We searched the following computerized databases: MEDLINE (1966 to December 2009), Scopus (1960 to December 2009), Web of Science (1975 to December 2009) and the Cochrane Library (1993 to 2009 [issue 4]). The complete MEDLINE search strategy is presented in Appendix 1 (available at www.cmaj.ca/cgi/content/full/cmaj.091786/DC1). In an attempt to minimize the omission of potentially relevant trials, we also searched the reference lists of included studies. We used the terms “patient care team,” “disease management,” “case management,” “managed care programs,” “home-based intervention” and “patient care management” to cover the inconsistency in the definition of disease management; we also used the terms “diabetes mellitus,” “HbA_{1c}” and “glycated hemoglobin.”

Two of us (C.P. and C.H.) first reviewed the titles and abstracts of identified articles and then examined the full-text version of selected articles further to assess relevance to the research topic. Only RCTs were included, because this study design supports maximum validity and causal

inference.⁹ The search was limited to English-language publications. We restricted inclusion to studies that reported hemoglobin A_{1c} levels, which is an index of the mean blood glucose concentration of the preceding 8–12 weeks.¹⁰ In addition to our operational definition of disease management, we defined the following inclusion criteria: the study had to involve adults with type 1 or 2 diabetes; it had to report both pre- and postintervention hemoglobin A_{1c} levels; and postintervention hemoglobin A_{1c} levels had to be assessed after at least 12 weeks of follow-up.

We excluded trials in which the intervention did not involve direct contact between the disease manager and the patient or was unclear, unspecified or exclusively based on contact by Internet or mail.

Data extraction

Two of us (C.P. and C.H.) evaluated each study separately and extracted data. To assess outcome, hemoglobin A_{1c} levels before and after the intervention were extracted. In the event of several postintervention values, only the first one was considered. Other data extracted were as follows: characteristics of the participants (percentage of women, mean age), sample size, number of dropouts, intervention mode (one-to-one session, phone contact or both), type of program components (patient education, psychological coaching, monitoring, feedback of initial evaluation to primary care physician, treatment adjustment), length of intervention, frequency of contact, interval between pre- and postintervention hemoglobin A_{1c} assessments, and adverse events (hypoglycemic episodes, hospital admission and death).

Frequency of contact was estimated on the basis of the reported intervention protocol and, when available, the results. We classified the frequency into three levels: low (less than one contact monthly per patient), moderate (one contact monthly per patient) and high (several contacts monthly per patient). In the event of discrepancies in the classification of contact frequency, data were reviewed by another one of us (M.L.G.), and a consensus was reached.

When data were missing, the original authors of the article were contacted by email.

Statistical analysis

To account for differences in baseline hemoglobin A_{1c} levels between the studies, we calculated the mean difference between pre- and postintervention hemoglobin A_{1c} levels for the intervention and control groups, and the standard deviation (SD) of each difference. Thus, our outcome corresponds to the improvement in glycemic control in the intervention group between base-

line and postintervention hemoglobin A_{1c} levels compared with the control group. We used the imputation method according to baseline values for missing SDs (we imputed missing SDs according to the pre-intervention values). Owing to significant heterogeneity, we used a random-effects model to calculate the pooled standardized mean difference in hemoglobin A_{1c} levels between the intervention and control groups, along with the 95% confidence interval (CI).¹¹ Heterogeneity was quantified by using *I*² and τ^2 (study variance) values.^{12,13}

We used meta-regression analysis to determine what part of between-study variance was explained by patient characteristics (mean hemoglobin A_{1c} level, age, sex) and components of the disease-management programs (length of intervention, treatment adjustment, mode of patient education, frequency of contact, feedback of initial evaluation to primary care physician, and mode of intervention). Results are expressed as standardized mean changes in the hemoglobin A_{1c} level. Explained heterogeneity was expressed as a percentage change of τ^2 (between-study variance).

Because quality assessment in meta-analysis is controversial,¹⁴ we performed three sensitivity analyses based on key components of internal validity to test the robustness of our results.¹⁵ In the first sensitivity analysis, we excluded trials that had a dropout rate of 20% or more and trials without dropout information. In the second analysis, we excluded trials in which the difference in dropout rates between study groups was 7% or more (highest quintile) and trials without dropout information. In the third analysis, we excluded trials with unclear information about allocation concealment.¹⁶

For all analyses, a *p* value of 0.05 or less was considered to be statistically significant.

Results

Study characteristics

The selection of studies for our review is summarized in Figure 1.¹⁷ The initial search strategy identified 2148 citations, and 135 full-text articles were reviewed. Forty-four studies met our inclusion criteria. Three studies were excluded because of missing data on hemoglobin A_{1c} levels at baseline, even after contacting the authors.^{18–20} Thus, we included 41 RCTs published between 1990 and 2009 that enrolled a total of 7013 adults with type 1 or 2 diabetes.^{21–61}

The main features of the 41 RCTs are shown in Table 1 (at the end of the article). Twenty-six trials were conducted in the United States, five in Canada, three in Europe and seven in Asia. Sam-

ple sizes ranged from 31 to 1665. The length of the intervention ranged from 1.5 to 48 months. In most trials, the length of intervention and the length of follow-up were similar, with only five trials reporting a few months' difference between the end of the intervention and hemoglobin A_{1c} assessment.^{21,23,29,39,49} Most of the studies (29 trials) focused solely on type 2 diabetes, 9 included patients with either type 1 or 2 diabetes, and 3 trials focused on type 1 diabetes. The mean age of the participants was 57.6 years (SD 7.3); 46.0% were men. The mean hemoglobin A_{1c} concentration at baseline was 8.5% (SD 1.4%).

Effect of intervention on glycemic control

The impact of the disease-management programs on changes in hemoglobin A_{1c} concentrations in the intervention and control groups is presented in Figure 2. In the random-effects model, the pooled standardized mean difference in levels between the intervention and control groups was -0.38 (95% CI -0.47 to -0.29 ; *p* < 0.001), favouring disease




Figure 1: Selection of randomized controlled trials (RCTs) for the meta-analysis.

management over usual care. This standardized mean difference corresponds to an absolute mean difference in hemoglobin A_{1c} levels of 0.51% between the intervention and control groups. None of the studies reported a significant change in hemoglobin A_{1c} in favour of usual care. There was significant heterogeneity among the trials regarding changes in hemoglobin A_{1c} ($I^2 = 66\%$).⁶²

Meta-regression and subgroup analyses

Results of univariable meta-regression analyses, stratified by patient characteristics and components of the disease-management programs, are shown in Table 2. Of the patient characteristics analyzed, age and sex were not associated with between-group differences in hemoglobin A_{1c} outcomes. The reduction in hemoglobin A_{1c} lev-




Figure 2: Estimated differences in hemoglobin A_{1c} level before and after intervention of disease management for improved glycemic control in adults with type 1 or 2 diabetes mellitus. Standardized mean differences between intervention and control groups of less than zero indicate an effect in favour of disease-management programs. CI = confidence interval, CMD study = California Medi-Cal Type 2 Diabetes Study.

els was significantly greater among patients with a baseline hemoglobin A_{1c} level of 8.0% or higher (standardized mean difference -0.45) than among those with a baseline level of less than 8.0% (standardized mean difference -0.14) ($p = 0.003$). About 33% of the variance between trials could be explained by mean hemoglobin A_{1c} values at baseline.

Two components of the disease-management programs led to greater improvements in glycaemic control (Table 2). First, programs in which the disease manager was able to start or modify

treatment with or without prior approval from the primary care physician resulted in a significantly greater reduction in hemoglobin A_{1c} levels (standardized mean difference -0.60 v. -0.28 in trials with no approval to do so; $p < 0.001$). Second, among the 36 trials that reported sufficient information to allow classification of the frequency of patient contact (Table 1), programs with a moderate or high frequency of contact (28 trials) reported a significant reduction in hemoglobin A_{1c} levels compared with usual care (standardized mean difference -0.56 for high frequency and

Table 2: Effect of patient characteristics and components of disease-management programs on changes in hemoglobin A_{1c} concentration

Variable	No. of studies	Standardized mean difference in change of hemoglobin A _{1c} between intervention and control groups (95% CI)	<i>p</i> value for difference in effect*	Heterogeneity, † <i>I</i> ² (95% CI), %	Variance between studies explained by variable, %
Patient characteristic					
Mean HbA _{1c} level at baseline	40‡				32.7
< 8.0%	11	-0.14 (-0.25 to -0.05)		25 (0 to 63)	
≥ 8.0%	29	-0.45 (-0.56 to -0.34)	0.003	59 (38 to 73)	
Age	36‡	0.01 (-0.01 to 0.02)	0.23		22.2
Sex	39‡	0.00 (-0.01 to 0.01)	0.30		4.6
Component of disease-management program					
Treatment adjustment	41				39.2
No (ref)	28	-0.28 (-0.37 to -0.18)		60 (39 to 73)	
Yes	13	-0.60 (-0.73 to -0.47)	< 0.001	28 (0 to 63)	
Patient education	39‡				31.9
Individual (ref)	31	-0.32 (-0.41 to -0.23)		54 (31 to 69)	
Group + individual	8	-0.48 (-0.68 to -0.28)	0.11	65 (25 to 84)	
Intervention mode	41				10.7
Phone (ref)	10	-0.27 (-0.42 to -0.12)		53 (5 to 77)	
Face to face + phone	18	-0.47 (-0.63 to -0.32)	0.12	71 (54 to 82)	
Face to face	13	-0.30 (-0.43 to -0.16)	0.90	39 (0 to 68)	
Length of intervention, mo	41				7.5
< 12 (ref)	19	-0.48 (-0.63 to -0.33)		53 (21 to 72)	
≥ 12	22	-0.31 (-0.42 to -0.20)	0.08	69 (52 to 80)	
Frequency of contact	36‡				6.1
Low (ref)	8	-0.30 (-0.54 to 0.06)		80 (62 to 90)	
Moderate	12	-0.24 (-0.37 to -0.12)	0.73	33 (0 to 66)	
High	16	-0.56 (-0.72 to -0.40)	0.033	52 (14 to 73)	
Feedback of initial evaluation to primary care physician	41				3.4
Yes (ref)	21	-0.33 (-0.44 to -0.22)		67 (49 to 79)	
No	20	-0.44 (-0.59 to -0.29)	0.26	58 (31 to 74)	

Note: CI = confidence interval, ref = reference group.

**p* values refer to meta-regression analysis. For each variable, the *p* value compares the effect of each category compared with the reference category.

†Values of < 50% represent a low level of heterogeneity, ≥ 50% to < 75% a moderate level of heterogeneity and ≥ 75% a high level of heterogeneity.

‡Number of trials does not total 41 because trials with missing data for the variable specified were excluded.

Table 3: Components of quality assessment that were considered for the sensitivity analyses

Study	Dropout rate, %	Difference in dropout rates between study groups, %	Allocation concealment
Estey et al. ²¹	11.7	NA	B
Ahring et al. ²²	9.5	2.1	B
Franz et al. ²³	27.5	NA	B
Weinberger et al. ²⁴	8.7	3.4	B
Jaber et al. ²⁵	13.3	26.1	B
Fukuda et al. ²⁶	3.8	3.0	B
Gaede et al. ²⁷	6.9	3.7	A
Ridgeway et al. ²⁸	32.1	7.1	B
Sadur et al. ²⁹	15.7	4.4	B
Thompson et al. ³⁰	0	0	A
Piette et al. ³¹	11.4	3.8	A
Hiss et al. ³²	27.4	3.7	B
Piette et al. ³³	6.2	5.5	A
Litaker et al. ³⁴	NA	NA	B
Oh et al. ³⁵	24.0	8.0	B
Polonsky et al. ³⁶	39.8	13.5	B
Taylor et al. ³⁷	24.8	5.0	B
Gary et al. ³⁸	23.9	1.0	A
Goudswaard et al. ³⁹	13.8	1.0	A
CMD Study ⁴⁰	12.4	7.0	A
Ko et al. ⁴¹	1.1	2.2	B
Krein et al. ⁴²	15.0	2.3	B
Montori et al. ⁴³	9.7	7.1	A
Choe et al. ⁴⁴	18.7	13.4	B
Farmer et al. ⁴⁵	12.9	8.9	A
Ménard et al. ⁴⁶	4.2	2.8	A
McMahon et al. ⁴⁷	19.2	7.7	A
Rothman et al. ⁴⁸	10.6	2.1	A
Taylor et al. ⁴⁹	2.5	0.05	B
Gabbay et al. ⁵⁰	NA	NA	B
Scott et al. ⁵¹	12.1	12.8	B
Shea et al. ⁵²	14.9	4.4	B
Shibayama et al. ⁵³	10.4	3.0	B
Hiss et al. ⁵⁴	16.7	3.9	B
Ko et al. ⁵⁵	29.5	5.2	A
Wattana et al. ⁵⁶	6.4	2.6	B
Sun et al. ⁵⁷	2.7	0.0	B
Dale et al. ⁵⁸	12.8	4.6	A
Doucette et al. ⁵⁹	15.4	2.8	B
Kim et al. ⁶⁰	4.8	2.3	B
Samuel-Hodge et al. ⁶¹	15.4	4.2	A

Note: A = adequate, B = unclear, CMD Study = California Medi-Cal Type 2 Diabetes Study, NA = not available.

−0.24 for moderate frequency). Nevertheless, only programs with a high frequency of contact led to a significantly greater reduction in hemoglobin A_{1c} levels compared with programs with a low frequency of contact (standardized mean difference −0.56 v. −0.30, $p = 0.03$).

In the random-effects subgroup analyses, none of the other program components modified the effectiveness of the intervention on hemoglobin A_{1c} levels. Two components explained a large part of the variance between trials: 31.9% was explained by mode of education and 39.2% by treatment adjustment. Trials in which the disease manager was able to start or modify treatment with or without prior approval of the physician, trials with face-to-face sessions and trials with a moderate frequency of patient contact each showed a low level of heterogeneity ($I^2 < 50\%$).

Sensitivity analyses

Our primary findings did not change after we excluded trials with dropout rates of 20% or more and trials without dropout information (Table 3). The same was true after we excluded trials with a between-group difference in dropout rates of 7% or more and trials without dropout information, and after we excluded trials with unclear allocation concealment (Table 3).

Adverse events

Hypoglycemic episodes were not systematically assessed. Only 9 of the 41 studies reported this information separately for intervention and control groups, but with varied definitions of hypoglycemic episodes.^{22,25,27,31,40,43,45–47} No difference in hypoglycemic episodes between study groups was reported in six of the nine trials. In two of the three trials that reported a difference, the adverse event occurred more frequently in the control groups than in the intervention groups.^{31,45} Twenty studies reported deaths over the follow-up period; no overall difference in mortality between groups was found ($p = 0.18$). Hospital admissions were not clearly or systematically reported.

Interpretation

Our meta-analysis suggests that disease-management programs have a favourable effect on improving glycemic control, with a pooled standardized mean reduction of 0.38 (corresponding to a pooled absolute mean reduction of 0.51%) in hemoglobin A_{1c} levels compared with usual care. This finding was robust in sensitivity analyses based on quality assessment. The United Kingdom Prospective Diabetes Study showed that each 1% reduction in hemoglobin A_{1c} level was associated with a 37% decrease in the risk of

microvascular complications and a 21% decrease in the risk of death related to diabetes, with no evidence of a threshold.⁶³ Therefore, the absolute reduction of 0.51% in hemoglobin A_{1c} level in our study appears to be clinically significant. Moreover, this finding is probably largely underestimated, because the usual care provided in control groups in RCTs is often better than that provided in clinical practice. Indeed, there was a significant standardized mean reduction in hemoglobin A_{1c} levels of -0.25 in the control groups, which corresponds to an absolute mean reduction of 0.40%. Some studies included in our meta-analysis permitted patients in the control group to contact the medical team or be contacted by them during follow-up in addition to usual care.^{23,43,55} Also, patients received structured individual education before randomization in some trials.^{21,23}

Our findings suggest that disease-management programs are more effective for patients who have poor glycemic control (mean hemoglobin A_{1c} \geq 8.0% at baseline) than for those with better glycemic control. This is concordant with results among patients starting insulin therapy.⁶⁴ Thus, disease management could be particularly effective if targeted at patients with nonstabilized diabetes. Moreover, such patients have a higher risk of complications and so would probably derive greater long-term benefit from disease management.

We found that the ability of disease managers to start or modify medical treatment was an effective feature of disease-management programs. This confirms the findings of Shojania and colleagues, who evaluated the ability to adjust treatment without prior physician agreement.⁷ However, we found that the ability to adjust treatment was an effective feature both with and without prior physician agreement, which is more relevant for physicians, nurses and pharmacists in clinical practice. This has important implications, because nonadherence to medical treatment is a significant predictor of all-cause mortality and hospital admission among patients with diabetes.⁶⁵

Despite its relevance for clinicians and policy-makers, the intensity of disease-management programs has not been investigated in previous reviews. Program intensity depends on the frequency of patient contacts, their duration and the length of the program. Because the duration of contact was not reported in most of the studies included in our review, we were not able to explore it. However, we explored frequency of contact and length of intervention. We did not find any significant difference associated with length of intervention, despite a nonsignificant improvement observed with shorter interventions. Frequency of contact proved to be a key feature of

the effectiveness of disease-management programs. There was substantial discrepancy in frequency across trials, ranging from “counseling by telephone every week if necessary”²² to “at least five visits by the nurse within a study period of one year.”⁴¹ For our analysis, frequency of contact was estimated on the basis of the intervention protocol reported and, when available, the results. Although the reported intervention protocol probably overestimated the real frequency of contact, frequency was evaluated on the basis of results in 12 studies and was consequently found to be an effective measure. Our findings are consistent with those from a recent large controlled trial, although it showed a nonsignificant trend toward better glycemic control with more intensive intervention.²⁰ The greater effectiveness associated with a high frequency of patient contact suggests that only disease-management programs with intensive interventions should be implemented, perhaps by targeting patients at high risk of diabetes complications.

Patient education is the cornerstone of diabetes care. An overall beneficial effect of education among patients with diabetes has already been shown in several studies.^{66,67} We did not find any difference in effectiveness between individual education and a combination of individual and group education. This finding suggests that a combination of group and individual education could be a solution to cope with the lack of medical providers and the time-consuming aspect of individual education. Surprisingly, neither the mode of contact nor feedback of the initial evaluation to the primary care physician were discriminatory components. However, we cannot rule out the possibility of incorrect classification of feedback as a program component, because it was taken for granted that such feedback would be provided systematically, so this step was not stipulated formally in the protocol.

Strengths and limitations

The strengths of the study include a comprehensive systematic review of the literature, with a large number of studies included. We used a broad search strategy to capture all relevant information. Our work confirms the findings of previous reviews, with a mean difference in hemoglobin A_{1c} level similar to that observed in previous studies.^{7,8,68,69} However, we included only RCTs and several more recent studies, with thus a larger sample size. Therefore, our estimate is probably more precise than that in previous studies.

Our study has limitations. Our analyses were based on results from randomized controlled trials, and adjustment was not done at an individual patient level. By including only studies published

in English, we may have missed other relevant studies. The weak description of the intervention strategy in most studies precluded the analysis of some potentially relevant components. Notably, we were unable to study the effect of the degree of the primary care physician's involvement in these programs, which is an essential aspect for implementation. For some components, such as frequency of patient contact, we contacted the authors for more details. However, because some trials were performed several years ago, no supplementary information was available. Another limitation was the short follow-up in many of the trials, even though we excluded trials with less than 12 weeks of follow-up. Because only five trials continued for more than 12 months, we were unable to capture the long-term effects of disease-management programs. However, outcomes such as long-term diabetes complications, especially vascular complications, have not yet been examined in studies of disease management for improved diabetes care. In some trials, the length of the intervention was very short (less than six months in six trials) and thus may have been too short to produce any clinical benefits.

We noted heterogeneity in the overall effect estimate and performed a meta-regression analysis to determine potential sources. The two components of disease-management programs that led to significantly greater improvements in glycemic control accounted for 6.1% (frequency of contact) and 39.2% (treatment adjustment) of the variance between studies. We did not identify all sources of variance among trials, but a meta-analysis of summary data from reported studies has little capacity to do so.

Although a recurrent problem in meta-analyses is publication bias, application of asymmetry tests seemed inappropriate owing to the presence of heterogeneity.⁷⁰ A previous meta-analysis reported a larger effect estimate for small studies.⁷ Because a higher intensity of intervention appears to be an important feature underpinning the efficacy of disease-management programs, this "size trial effect" could be due to a higher intensity of intervention in small studies. Indeed, of the 16 studies with a high frequency of patient contact in our analysis, 11 (69%) were relatively small, with samples smaller than the median for the studies included (117 patients). This more intensive intervention in small studies, rather than publication bias, could explain the greater improvement in glycemic control.

Conclusion

Disease-management programs had a clinically moderate but significant impact on hemoglobin A_{1c} levels among adults with diabetes. Effective

components of the programs were a high frequency of patient contact and the ability for disease managers to adjust treatment with or without prior physician approval. Our findings have important implications for both the current policy on the delivery of diabetes care and the direction of future research. Our work delineates a general framework with core features for effective programs for disease management. Priority should be given to programs with intensive and proactive follow-up that target patients at high risk of diabetes complications rather than to programs with low frequency of contact that target the overall population of patients with diabetes. In addition, disease managers should be allowed to start or modify medical treatment proactively.

More research is needed concerning the long-term impact of disease-management programs on glycemic control, microvascular and macrovascular complications, admission to hospital and mortality. Further research should also determine whether, in addition to patients with nonstabilized diabetes, other groups of patients with diabetes would benefit from disease management. Lastly, high-quality cost-effectiveness studies of disease-management programs are needed to direct care providers and policy-makers in the allocation of health care resources.

References

1. Global Guideline for Type 2 Diabetes: recommendations for standard, comprehensive, and minimal care. *Diabet Med* 2006;23:579-93.
2. Feder G, Griffiths C, Highton C, et al. Do clinical guidelines introduced with practice based education improve care of asthmatic and diabetic patients? A randomised controlled trial in general practices in east London. *BMJ* 1995;311:1473-8.
3. Davis DA, Taylor-Vaisey A. Translating guidelines into practice. A systematic review of theoretic concepts, practical experience and research evidence in the adoption of clinical practice guidelines. *CMAJ* 1997;157:408-16.
4. Coster S, Gulliford MC, Seed PT, et al. Self-monitoring in type 2 diabetes mellitus: a meta-analysis. *Diabet Med* 2000;17:755-61.
5. Vermeire E, Wens J, Van Royen P, et al. Interventions for improving adherence to treatment recommendations in people with type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2005;CD003638.
6. Farmer A, Gibson OJ, Tarassenko L, et al. A systematic review of telemedicine interventions to support blood glucose self-monitoring in diabetes. *Diabet Med* 2005;22:1372-8.
7. Shojania KG, Ranji SR, McDonald KM, et al. Effects of quality improvement strategies for type 2 diabetes on glycemic control: a meta-regression analysis. *JAMA* 2006;296:427-40.
8. Chodosh J, Morton SC, Mojica W, et al. Meta-analysis: chronic disease self-management programs for older adults. *Ann Intern Med* 2005;143:427-38.
9. Richter B, Berger M. Randomized controlled trials remain fundamental to clinical decision making in Type II diabetes mellitus: a comment to the debate on randomized controlled trials (For debate). *Diabetologia* 2000;43:528-32.
10. Nathan DM, Turgeon H, Regan S. Relationship between glycated haemoglobin levels and mean glucose levels over time. *Diabetologia* 2007;50:2239-44.
11. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177-88.
12. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539-58.
13. Rucker G, Schwarzer G, Carpenter JR, et al. Undue reliance on I(2) in assessing heterogeneity may mislead. *BMC Med Res Methodol* 2008;8:79.

14. Jüni P, Witschi A, Bloch R, et al. The hazards of scoring the quality of clinical trials for meta-analysis. *JAMA* 1999;282:1054-60.
15. Jüni P, Altman DG, Egger M. Systematic reviews in health care: Assessing the quality of controlled clinical trials. *BMJ* 2001;323:42-6.
16. Ryan R, Hill S, Broclain D, et al. *Cochrane Consumers and Communication Review Group: study quality guide*. Victoria (Australia); 2007.
17. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Ann Intern Med* 2009;151:W65-94.
18. Aubert RE, Herman WH, Waters J, et al. Nurse case management to improve glycemic control in diabetic patients in a health maintenance organization. A randomized, controlled trial. *Ann Intern Med* 1998;129:605-12.
19. Groeneveld Y, Petri H, Hermans J, et al. An assessment of structured care assistance in the management of patients with type 2 diabetes in general practice. *Scand J Prim Health Care* 2001;19:25-30.
20. Gary TL, Batts-Turner M, Yeh HC. The effects of a nurse case manager and a community health worker team on diabetic control, emergency department visits, and hospitalizations among urban African Americans with type 2 diabetes mellitus: a randomized controlled trial. *Arch Intern Med* 2009;169:1788-94.
21. Estey AL, Tan MH, Mann K. Follow-up intervention: its effect on compliance behavior to a diabetes regimen. *Diabetes Educ* 1990;16:291-5.
22. Ahring KK, Ahring JP, Joyce C, et al. Telephone modem access improves diabetes control in those with insulin-requiring diabetes. *Diabetes Care* 1992;15:971-5.
23. Franz MJ, Monk A, Barry B, et al. Effectiveness of medical nutrition therapy provided by dietitians in the management of non-insulin-dependent diabetes mellitus: a randomized, controlled clinical trial. *J Am Diet Assoc* 1995;95:1009-17.
24. Weinberger M, Kirkman MS, Samsa GP, et al. A nurse-coordinated intervention for primary care patients with non-insulin-dependent diabetes mellitus: impact on glycemic control and health-related quality of life. *J Gen Intern Med* 1995;10:59-66.
25. Jaber LA, Halapy H, Fernet M, et al. Evaluation of a pharmaceutical care model on diabetes management. *Ann Pharmacother* 1996;30:238-43.
26. Fukuda H, Muto T, Kawamori R. Evaluation of a diabetes patient education program consisting of a three-day hospitalization and a six-month follow-up by telephone counseling for mild diabetes and IGT. *Environ Health Prev Med* 1999;4:122-9.
27. Gaede P, Vedel P, Parving HH, et al. Intensified multifactorial intervention in patients with type 2 diabetes mellitus and microalbuminuria: the Steno type 2 randomised study. *Lancet* 1999;353:617-22.
28. Ridgeway NA, Harvill DR, Harvill LM, et al. Improved control of type 2 diabetes mellitus: a practical education/behavior modification program in a primary care clinic. *South Med J* 1999;92:667-72.
29. Sadur CN, Moline N, Costa M, et al. Diabetes management in a health maintenance organization. Efficacy of care management using cluster visits. *Diabetes Care* 1999;22:2011-7.
30. Thompson DM, Kozak SE, Sheps S. Insulin adjustment by a diabetes nurse educator improves glucose control in insulin-requiring diabetic patients: a randomized trial. *CMAJ* 1999;161:959-62.
31. Piette JD, Weinberger M, McPhee SJ, et al. Do automated calls with nurse follow-up improve self-care and glycemic control among vulnerable patients with diabetes? *Am J Med* 2000;108:20-7.
32. Hiss RG, Gillard ML, Armbruster BA, et al. Comprehensive evaluation of community-based diabetic patients: effect of feedback to patients and their physicians: a randomized controlled trial. *Diabetes Care* 2001;24:690-4.
33. Piette JD, Weinberger M, Kraemer FB, et al. Impact of automated calls with nurse follow-up on diabetes treatment outcomes in a Department of Veterans Affairs Health Care System: a randomized controlled trial. *Diabetes Care* 2001;24:202-8.
34. Litaker D, Mion L, Planavsky L, et al. Physician-nurse practitioner teams in chronic disease management: the impact on costs, clinical effectiveness, and patients' perception of care. *J Interprof Care* 2003;17:223-37.
35. Oh JA, Kim HS, Yoon KH, et al. A telephone-delivered intervention to improve glycemic control in type 2 diabetic patients. *Yonsei Med J* 2003;44:1-8.
36. Polonsky WH, Earles J, Smith S, et al. Integrating medical management with diabetes self-management training: a randomized control trial of the Diabetes Outpatient Intensive Treatment program. *Diabetes Care* 2003;26:3048-53.
37. Taylor CB, Miller NH, Reilly KR, et al. Evaluation of a nurse-care management system to improve outcomes in patients with complicated diabetes. *Diabetes Care* 2003;26:1058-63.
38. Gary TL, Bone LR, Hill MN, et al. Randomized controlled trial of the effects of nurse case manager and community health worker interventions on risk factors for diabetes-related complications in urban African Americans. *Prev Med* 2003;37:23-32.
39. Goudswaard AN, Stolk RP, Zuihthoff NP, et al. Long-term effects of self-management education for patients with type 2 diabetes taking maximal oral hypoglycaemic therapy: a randomized trial in primary care. *Diabet Med* 2004;21:491-6.
40. California Medi-Cal Type 2 Diabetes Study Group. Closing the gap: effect of diabetes case management on glycemic control among low-income ethnic minority populations: the California Medi-Cal Type 2 Diabetes Study. *Diabetes Care* 2004;27:95-103.
41. Ko GT, Li JK, Kan EC, et al. Effects of a structured health education programme by a diabetic education nurse on cardiovascular risk factors in Chinese type 2 diabetic patients: a 1-year prospective randomized study. *Diabet Med* 2004;21:1274-9.
42. Krein SL, Klamers ML, Vijan S, et al. Case management for patients with poorly controlled diabetes: a randomized trial. *Am J Med* 2004;116:732-9.
43. Montori VM, Helgemoe PK, Guyatt GH, et al. Telecare for patients with type 1 diabetes and inadequate glycemic control: a randomized controlled trial and meta-analysis. *Diabetes Care* 2004;27:1088-94.
44. Choe HM, Mitrovich S, Dubay D, et al. Proactive case management of high-risk patients with type 2 diabetes mellitus by a clinical pharmacist: a randomized controlled trial. *Am J Manag Care* 2005;11:253-60.
45. Farmer AJ, Gibson OJ, Dudley C, et al. A randomized controlled trial of the effect of real-time telemedicine support on glycemic control in young adults with type 1 diabetes. *Diabetes Care* 2005;28:2697-702.
46. Menard J, Payette H, Baillargeon JP, et al. Efficacy of intensive multitherapy for patients with type 2 diabetes mellitus: a randomized controlled trial. *CMAJ* 2005;173:1457-66.
47. McMahon GT, Gomes HE, Hickson Hohne S, et al. Web-based care management in patients with poorly controlled diabetes. *Diabetes Care* 2005;28:1624-9.
48. Rothman RL, Malone R, Bryant B, et al. A randomized trial of a primary care-based disease management program to improve cardiovascular risk factors and glycated hemoglobin levels in patients with diabetes. *Am J Med* 2005;118:276-84.
49. Taylor KI, Oberle KM, Crutcher RA, et al. Promoting health in type 2 diabetes: nurse-physician collaboration in primary care. *Biol Res Nurs* 2005;6:207-15.
50. Gabbay RA, Lendel I, Saleem TM, et al. Nurse case management improves blood pressure, emotional distress and diabetes complication screening. *Diabetes Res Clin Pract* 2006;71:28-35.
51. Scott DM, Boyd ST, Stephan M, et al. Outcomes of pharmacist-managed diabetes care services in a community health center. *Am J Health Syst Pharm* 2006;63:2116-22.
52. Shea S, Weinstock RS, Starren J, et al. A randomized trial comparing telemedicine case management with usual care in older, ethnically diverse, medically underserved patients with diabetes mellitus. *J Am Med Assoc* 2006;296:1340-51.
53. Shibayama T, Kobayashi K, Takano A, et al. Effectiveness of lifestyle counseling by certified expert nurse of Japan for non-insulin-treated diabetic outpatients: a 1-year randomized controlled trial. *Diabetes Res Clin Pract* 2007;76:265-8.
54. Hiss RG, Armbruster BA, Gillard ML, et al. Nurse care manager collaboration with community-based physicians providing diabetes care: a randomized controlled trial. *Diabetes Educ* 2007;33:493-502.
55. Ko SH, Song KH, Kim SR, et al. Long-term effects of a structured intensive diabetes education programme (SIDEPE) in patients with type 2 diabetes mellitus — a 4-year follow-up study. *Diabet Med* 2007;24:55-62.
56. Wattana C, Srisuphan W, Pothiban L, et al. Effects of a diabetes self-management program on glycemic control, coronary heart disease risk, and quality of life among Thai patients with type 2 diabetes. *Nurs Health Sci* 2007;9:135-41.
57. Sun J, Wang Y, Chen X, et al. An integrated intervention program to control diabetes in overweight Chinese women and men with type 2 diabetes. *Asia Pac J Clin Nutr* 2008;17:514-24.
58. Dale J, Caramlau I, Sturt J, et al. Telephone peer-delivered intervention for diabetes motivation and support: the telecare exploratory RCT. *Patient Educ Couns* 2009;75:91-8.
59. Doucette WR, Witry MJ, Farris KB, et al. Community pharmacist-provided extended diabetes care. *Ann Pharmacother* 2009;43:882-9.
60. Kim MT, Han HR, Song HJ, et al. A community-based, culturally tailored behavioral intervention for Korean Americans with

- type 2 diabetes. *Diabetes Educ* 2009;35:986-94.
61. Samuel-Hodge CD, Keyserling TC, Park S, et al. A randomized trial of a church-based diabetes self-management program for African Americans with type 2 diabetes. *Diabetes Educ* 2009; 35:439-54.
 62. Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557-60.
 63. Stratton IM, Adler AI, Neil HA, et al. Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ* 2000;321:405-12.
 64. Hayward RA, Manning WG, Kaplan SH, et al. Starting insulin therapy in patients with type 2 diabetes: effectiveness, complications, and resource utilization. *JAMA* 1997;278:1663-9.
 65. Ho PM, Rumsfeld JS, Masoudi FA, et al. Effect of medication nonadherence on hospitalization and mortality among patients with diabetes mellitus. *Arch Intern Med* 2006;166:1836-41.
 66. Brown SA. Studies of educational interventions and outcomes in diabetic adults: a meta-analysis revisited. *Patient Educ Couns* 1990;16:189-215.
 67. Ellis SE, Speroff T, Dittus RS, et al. Diabetes patient education: a meta-analysis and meta-regression. *Patient Educ Couns* 2004; 52:97-105.
 68. Knight K, Badamgarav E, Henning JM, et al. A systematic review of diabetes disease management programs. *Am J Manag Care* 2005;11:242-50.
 69. Norris SL, Nichols PJ, Caspersen CJ, et al. The effectiveness of disease and case management for people with diabetes. A systematic review. *Am J Prev Med* 2002;22:15-38.
 70. Ioannidis JP, Trikalinos TA. The appropriateness of asymmetry tests for publication bias in meta-analyses: a large survey. *CMAJ* 2007;176:1091-6.

Competing interests: Catherine Helmer has received grants from the Agence nationale de la recherche and the Institut national de prévention et d'éducation pour la santé; honoraria

from Eisai Inc., Pfizer Inc. and Ipsen; and royalties from Edimark for an article on an unrelated topic. Rodolphe Thiébaud is a member of the advisory board for Gilead. No competing interests declared by Clément Pimouguet, Mélanie Le Goff or Jean François Dartigues.

Affiliations: From the Institut national de la santé et de la recherche médicale (Pimouguet, Le Goff, Thiébaud, Dartigues, Helmer); the Université Victor Segalen Bordeaux 2 (Pimouguet, Le Goff, Thiébaud, Dartigues, Helmer); and the Centre hospitalier universitaire de Bordeaux (Dartigues), Bordeaux, France

Contributors: Clément Pimouguet and Catherine Helmer designed the study, completed the literature search, extracted the data and drafted the manuscript. Mélanie Le Goff contributed to the data extraction, performed the data analysis and helped with the revision of the manuscript. Rodolphe Thiébaud helped with the interpretation of the results, and the consolidation and revision of the manuscript. Jean François Dartigues assisted with the design of the study, contributed to the interpretation of the results and helped with the consolidation and revision of the last draft of the manuscript. All of the authors approved the final version of the manuscript.

Funding: This study was funded by the French Ministry of Health. The ministry had no role in the design of the study, the analysis or interpretation of the data, or the writing of the manuscript.

Acknowledgements: The authors acknowledge the assistance of Evelyne Mouillet, Coralie Thore and Christian Weller from the documentary centre of the Institut de santé publique, d'épidémiologie et de développement.

Table 1: Details of randomized controlled trials of disease-management programs included in the meta-analysis (part 1 of 3)

Study (country)	No. of patients, intervention/control	Main inclusion criteria	Treatment in control group	Length of intervention, maximum no. of months	Modes of intervention	Frequency of contact	Team	Intervention component								
								Individual education	Group education	Coaching	Monitoring	PCP feedback information	Treatment adjustment			
Estey et al. ²¹ (Canada)	30/30	Type 2 diabetes	3-day education program	2.5	Face to face, phone	High	Nurse	X	X	X						
Ahring et al. ²² (Canada)	22/20	Type 1 diabetes; HbA _{1c} ≥ 7%; age 15–65 yr	Usual care	3	Phone	High	NA	X	X	X						X
Franz et al. ²³ (USA)	94/85	Type 2 diabetes; age 38–76 yr	One visit with dietitian	1.5	Face to face	Moderate	Dietitian	X	X	X						X
Weinberger et al. ²⁴ (USA)	204/71	Type 2 diabetes; age at onset of diabetes ≥ 40 yr	Usual care	12	Phone	Moderate	Nurse	X	X	X						X
Jaber et al. ²⁵ (USA)	23/22	Type 2 diabetes; African-American people	Usual care	4	Face to face	High	Pharmacist	X	X	X						X
Fukuda et al. ²⁶ (Japan)	27/25	Type 2 diabetes or IGT but no drug therapy for diabetes; HbA _{1c} < 8%; age 40–70 yr	Usual care	6	Face to face, phone	Moderate	Nurse	X	X	X						X
Gaede et al. ²⁷ (Denmark)	80/80	Type 2 diabetes; AER ≥ 30 mg/24 h; age 40–65 yr	Usual care	48	Face to face	NA	Nurse, dietitian, doctor	X	X	X						X
Ridgeway et al. ²⁸ (USA)	28/28	Type 2 diabetes; age ≥ 15–65 yr; ≥ 20% over ideal weight; fasting blood glucose > 150 mg/dL; HbA _{1c} > 6%	Usual care	12	Face to face	Moderate	Nurse, dietitian	X	X	X						
Sadur et al. ²⁹ (USA)	97/88	Type 1 or 2 diabetes; age 16–75 yr; HbA _{1c} > 8.5% or no HbA _{1c} measure during previous yr	Usual care	6	Face to face, phone	High	Nurse, dietitian, diabetologist, behaviourist, pharmacist	X	X	X						X
Thompson et al. ³⁰ (Canada)	23/23	Type 1 or 2 diabetes; insulin treatment; HbA _{1c} > 8.5%	Usual care	6	Phone	High	Nurse	X	X	X						X
Piette et al. ³¹ (USA)	124/124	Type 1 or 2 diabetes; age < 75 yr	Usual care	12	Phone	High	Nurse	X	X	X						X
Hiss et al. ³² (USA)	186/190	Type 2 diabetes	Usual care	12	Face to face	Low	Nurse	X	X	X						X
Piette et al. ³³ (USA)	146/146	Type 1 or 2 diabetes; age < 75 yr	Usual care	12	Phone	Moderate	Nurse	X	X	X						X
Litaker et al. ³⁴ (USA)	79/78	Type 2 diabetes; mild or moderate hypertension	Usual care	12	Face to face, phone	NA	Nurse	X	X	X						X
Oh et al. ³⁵ (South Korea)	25/25	Type 2 diabetes; HbA _{1c} ≥ 7%	Usual care	3	Phone	High	Dietitian, researcher	X	X	X						X

Table 1: Details of randomized controlled trials of disease-management programs included in the meta-analysis (part 2 of 3)

Study (country)	No. of patients, intervention/control	Main inclusion criteria	Treatment in control group	Length of intervention, maximum no. of months	Modes of intervention	Frequency of contact	Team	Intervention component					
								Individual education	Group education	Coaching	Monitoring information	PCP feedback	Treatment adjustment
Polonsky et al. ³⁶ (USA)	89/78	Type 1 or 2 diabetes; HbA _{1c} ≥ 8.5%	Quarterly educational mailings	6	Face to face, phone	Low	Nurse, physician, dietitian, exercise physiologist	X	X	X	X	X	X
Taylor et al. ³⁷ (USA)	84/85	Type 1 or 2 diabetes; HbA _{1c} > 10%	Usual care	12	Face to face, phone	Moderate	Nurse	X	X	X	X	X	X
Gary et al. ³⁸ (USA)	NA	Type 2 diabetes; age 35–75 yr	Informational mailings every 3–4 mo	24	Face to face, phone	Low	Nurse, community health worker	X	X	X	X	X	X
Goudswaard et al. ³⁹ (Netherlands)	28/30	Type 2 diabetes; HbA _{1c} ≥ 7%; age ≤ 75 yr	Usual care	6	Face to face	Moderate	Nurse	X	X	X	X	X	X
CMD Study ⁴⁰ (USA)	186/172	Type 2 diabetes; HbA _{1c} ≥ 7.5%	Usual care	36	Face to face, phone	NA	Nurse, dietitian	X	X	X	X	X	X
Ko et al. ⁴¹ (China)	90/88	Type 2 diabetes; HbA _{1c} 8%–11%; age 75–70 yr	Usual care	12	Face to face	Low	Nurse	X	X	X	X	X	X
Krein et al. ⁴² (USA)	123/123	Type 2 diabetes; HbA _{1c} ≥ 8.5%	Usual care	18	Face to face, phone	Low	Nurse	X	X	X	X	X	X
Montori et al. ⁴³ (USA)	15/16	Type 1 diabetes; HbA _{1c} ≥ 7.8%	Control patients could contact study nurse if necessary	6	Face to face, phone	High	Nurse, endocrinologist	X	X	X	X	X	X
Choe et al. ⁴⁴ (USA)	41/39	Type 2 diabetes; HbA _{1c} ≥ 8%; age ≤ 70 yr	Usual care	12	Face to face, phone	Moderate	Pharmacist	X	X	X	X	X	X
Farmer et al. ⁴⁵ (UK)	47/46	Type 1 diabetes; HbA _{1c} 8–11; age 18–30 yr	Minimal feedback of blood glucose	9	Phone	High	Nurse	X	X	X	X	X	X
Ménard et al. ⁴⁶ (Canada)	36/36	Type 2 diabetes; HbA _{1c} ≥ 8%; age 30–70 yr	Usual care	12	Face to face, phone	High	NA	X	X	X	X	X	X
McMahon et al. ⁴⁷ (USA)	52/52	Type 1 or 2 diabetes; HbA _{1c} ≥ 9%	Usual care	12	Web, phone	NA	Nurse	X	X	X	X	X	X
Rotlman et al. ⁴⁸ (USA)	112/105	Type 2 diabetes; HbA _{1c} ≥ 8%	Usual care	12	Face to face, phone	High	Pharmacist, diabetes care coordinator	X	X	X	X	X	X
Taylor et al. ⁴⁹ (Canada)	20/19	Type 2 diabetes	Usual care	3	Face to face	High	Nurse, dietitian, sport coach	X	X	X	X	X	X
Gabbay et al. ⁵⁰ (USA)	150/182	Type 1 or 2 diabetes; age ≥ 18 yr	Usual care	12	Face to face, phone	Low	Nurse	X	X	X	X	X	X

Table 1: Details of randomized controlled trials of disease-management programs included in the meta-analysis (part 3 of 3)

Study (country)	No. of patients, intervention/control	Main inclusion criteria	Treatment in control group	Length of intervention, maximum no. of months	Modes of intervention	Frequency of contact	Team	Intervention component					
								Individual education	Group education	Coaching	Monitoring	PCP feedback information	Treatment adjustment
Scott et al. ⁵¹ (USA)	76/73	Type 2 diabetes; age \geq 18 yr	Usual care	9	Face to face, phone	High	Nurse, pharmacist	X	X	X	X	X	X
Shea et al. ⁵² (USA)	844/821	Type 1 or 2 diabetes; age \geq 55 yr	Usual care	12	Phone	NA	Nurse, endocrinologist	X	X	X	X	X	X
Shibayama et al. ⁵³ (Japan)	67/67	Type 2 diabetes; age 20–75 yr; HbA _{1c} 6.5%–8.5%	Usual care	12	Face to face	Moderate	Nurse	X	X	X	X	X	X
Hiss et al. ⁵⁴ (USA)	95/102	Type 2 diabetes	Feedback of initial evaluation to primary care physician	6	Face to face	Moderate	Nurse	X	X	X	X	X	X
Ko et al. ⁵⁵ (South Korea)	219/218	Type 2 diabetes; age \leq 70 yr	Follow up every 3 mo; focused on blood glucose and drug adjustment	48	Face to face, phone	Low	Nurse, dietician, pharmacist, general practitioner, endocrinologist	X	X	X	X	X	X
Wattana et al. ⁵⁶ (Thailand)	75/72	Type 2 diabetes; age \geq 35 yr; fasting blood glucose $>$ 140 mg for \geq two follow-up visits	Usual care	6	Face to face	Moderate	Nurse	X	X	X	X	X	X
Sun et al. ⁵⁷ (China)	100/50	Type 2 diabetes; age 18–70 yr; BMI \geq 23 kg/m ²	Diet and physical instruction	6	Face to face	High	Dietician, physician	X	X	X	X	X	X
Dale et al. ⁵⁸ (United Kingdom)	44/97	Type 2 diabetes; HbA _{1c} $>$ 8%	Usual care	6	Phone	Moderate	Nurse	X	X	X	X	X	X
Doucette et al. ⁵⁹ (USA)	36/42	Type 2 diabetes; HbA _{1c} \geq 7%	Usual care	12	Face to face	Low	Pharmacist	X	X	X	X	X	X
Kim et al. ⁶⁰ (USA)	40/39	Type 2 diabetes; age \geq 30 yr; HbA _{1c} \geq 7.5%	Usual care	7.5	Face to face, phone	High	Nurse	X	X	X	X	X	X
Samuel-Hodge et al. ⁶¹ (USA)	117/84	Type 2 diabetes	Mailing to participants of 2 pamphlets and 3 monthly newsletters	12	Face to face, phone	High	Dietician, health professional, peer counsellor	X	X	X	X	X	X

Note: AER = urinary albumin excretion rate, BMI = body mass index, CMD Study = California Medi-Cal Type 2 Diabetes Study, IGT = impaired glucose tolerance, NA = not available, PCP = primary care physician.