CLINICAL IMAGES

Community-acquired pneumonia

Nicola Mumoli MD, Marco Cei MD

See also practice article by Leis and Gold on page 559 and at www.cmaj.ca/lookup/doi/10.1503/cmaj.111143**Competing interests:** None declared.

This article has been peer reviewed.

Affiliation: From the Department of Internal Medicine, Ospedale Civile Livorno, Livorno, Italy**Correspondence to:** Dr. Nicola Mumoli, n.mumoli@usl6.toscana.it

CMAJ 2012; DOI:10.1503/cmaj.111256

A 39-year-old man presented with fever (38.7°C), cough and right chest pain on inspiration. His blood pressure was 125/75 mm Hg, heart rate was 88 beats/min, respiration was 20 breaths/min and oxygen saturation on room air was 94%. Physical examination was remarkable only for crackles over the right upper lung field. Results of blood tests showed an elevated leukocyte count of 16.5 (normal 4.0 – 10.0) $\times 10^9/L$ and were otherwise normal. Urinary testing for *Legionella pneumophila* and *Streptococcus pneumoniae* antigens was negative. Chest radiography showed pneumonia in the right upper lobe (Figure 1). Because the patient was at low risk for death (class I on the Pneumonia Severity Index),¹ we prescribed oral penicillin and clarithromycin, and discharged the patient home. Follow-up chest radiography one month later showed that the consolidation had resolved (Appendix 1, available at www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.111256/-DC1).

About 4 million instances of community-acquired pneumonia occur in the United States each year. It is one of the most common causes

of death from infectious diseases worldwide.² The precise diagnosis continues to be complex; at least 50% of causative pathogens remain unidentified, and all rigorous definitions of pneumonia require the finding of a pulmonary infiltrate on chest radiography.² The initial antibiotic regimen should be chosen empirically to cover common typical and atypical pathogens. About 50% of patients with pneumonia can receive treatment outside of the hospital.^{2,3} The Pneumonia Severity Index is a validated prediction rule that identifies people with community-acquired pneumonia who are at low risk for death within 30 days of presentation.¹ An alternative scale is the CURB-65, which takes into account confusion, urea level, respiration, blood pressure and age.⁴ A general rule of thumb is to admit patients with oxygen saturation under 92%.⁵

References

1. Fine MJ, Auble TE, Yealy DM, et al. A prediction rule to identify low-risk patients with community-acquired pneumonia. *N Engl J Med* 1997;336:243–50.
2. Halm EA, Teirstein AS. Management of community-acquired pneumonia. *N Engl J Med* 2002;347:2039–45.
3. Mandell LA, Wunderink RG, Anzueto A, et al. Infectious Diseases Society of America/American Thoracic Society consensus guidelines on the management of community-acquired pneumonia in adults. *Clin Infect Dis* 2007;44(suppl 2):S27–72.
4. Chalmers JD, Singanayagam A, Akram AR, et al. Severity assessment tools for predicting mortality in hospitalised patients with community-acquired pneumonia. Systematic review and metaanalysis. *Thorax* 2010;65:878–83.
5. Majumdar SR, Eurich DT, Gamble JM, et al. Oxygen saturations less than 92% are associated with major adverse events in outpatients with pneumonia: a population-based cohort study. *Clin Infect Dis* 2011;52:325–31.

Helpful resources

- The Pneumonia Severity Index calculator is available online at <http://pda.ahrq.gov/clinic/psi/psicalc.asp>
- CURB-65 is available online at www.mdcalc.com/curb-65-severity-score-community-acquired-pneumonia

Figure 1: Posteroanterior and lateral radiographs of the chest of a 39-year-old man with fever, cough and pleuritic chest pain showing consolidation in the right upper lobe.