

Palliative care: renaming as supportive care and integration into comprehensive cancer care

Anthony J. Caprio MD

See also page E217 and www.cmaj.ca/lookup/doi/10.1503/cmaj.151171, and page 707 and www.cmaj.ca/lookup/doi/10.1503/cmaj.160547

Early palliative care for patients with advanced cancer was shown to improve quality of life, symptom management and satisfaction with care in three major randomized controlled trials (RCTs).¹⁻³ These findings prompted the inclusion of palliative care in clinical practice guidelines for patients with advanced cancer.^{4,5} Unfortunately, late referrals to palliative care and hospice services limit these potential benefits for patients and their caregivers.

In a linked study, Zimmermann and colleagues⁶ provide further evidence that negative perceptions of palliative care represent a potential barrier to early palliative care in the setting of advanced cancer. Using a sample of participants in a previous RCT of palliative care,³ the authors conducted qualitative interviews with patients and their caregivers from both the palliative care intervention group and standard oncology care control group.

This qualitative study reported several important findings. First, both groups described negative initial impressions of palliative care. However, the intervention group described a shift toward a more positive view through their experience of palliative care. Their experiences of palliative care practitioners partnering with patients, caregivers and oncologists to enhance quality of life and address physical and psychosocial distress reframed their concept of palliative care. Notably, the term “palliative care” was viewed as stigmatizing by both groups, whereas the alternative name “supportive care” was viewed more favourably by those patients and caregivers receiving palliative care.

The negative connotation of palliative care should not be surprising. Too often palliative care has been associated with death and dying, a loss of control or abandoning of hope. The term supportive care has generally been regarded more favourably. In a randomized trial of language used to describe palliative care services, the term supportive care was associated with more favourable impressions and a better understanding of services for patients with advanced

cancer.⁷ Similarly, oncology health care professionals preferred the name supportive care and considered the negative connotations associated with palliative care as a barrier to referral.⁸ In response to these findings, some palliative care programs chose to rebrand themselves as supportive care services.⁹

The interviews in the current study by Zimmerman and colleagues⁶ also show that patients and caregivers view health care professionals as having an important role in explaining and recommending palliative care. Although changing the name to supportive care may help promote a more positive view of palliative care, stigma will persist if this type of care is recommended only as default treatment when curative or life-prolonging treatments are deemed ineffective or undesired.

A shift in oncology care is necessary to facilitate early, concurrent palliative care to maximize outcomes for patients with cancer. Instead of dichotomizing standard oncology care and palliative care, palliative care should start at the time of cancer diagnosis and continue across the spectrum of cancer care. The type and intensity of palliative care services are calibrated over time to meet the needs of the patient and caregivers, regardless of prognosis or cancer outcome.^{4,10}

If palliative care is viewed as an essential evidence-based component of advanced disease management, then the benefits of palliative care support should be available and recommended to all patients and caregivers at every stage of their illness. Palliative care would no longer be chosen; it would simply be integrated into comprehensive

Competing interests: None declared.

This article was solicited and has not been peer reviewed.

Correspondence to:

Anthony Caprio,
Anthony.Caprio@
carolinashhealthcare.org

CMAJ 2016. DOI:10.1503/
cmaj.160206

KEY POINTS

- Early palliative care has been shown to improve quality of life, symptom management and satisfaction with care in patients with advanced cancer.
- Stigma associated with the term palliative care may be a barrier to timely referral and deprives patients and caregivers of the full benefits of palliative care.
- Although the term supportive care may be viewed more favourably, barriers to implementation of either supportive or palliative care can only be addressed by complete integration into routine oncology care.

oncology treatment that supports patients and caregivers by incorporating palliative care principles and services into all aspects of their care.

References

1. Bakitas M, Lyons KD, Hegel MT, et al. Effects of a palliative care intervention on clinical outcomes in patients with advanced cancer: the Project ENABLE II randomized controlled trial. *JAMA* 2009;302:741-9.
2. Temel JS, Greer JA, Muzikansky A, et al. Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010;363:733-42.
3. Zimmermann C, Swami N, Krzyzanowska M, et al. Early palliative care for patients with advanced cancer: a cluster-randomised controlled trial. *Lancet* 2014;383:1721-30.
4. Fassbender K, Watanabe SM. Early palliative care and its translation into oncology practice in Canada: barriers and challenges. *Ann Palliat Med* 2015;4:135-49.
5. Smith TJ, Temin S, Alesi ER, et al. American Society of Clinical Oncology provisional clinical opinion: the integration of palliative care into standard oncology care. *J Clin Oncol* 2012;30:880-7.
6. Zimmerman C, Swami N, Krzyzanowska M, et al. Perceptions of palliative care among patients with advanced cancer and their caregivers. *CMAJ* 2016;188:E217-27.
7. Maciasz RM, Arnold RM, Chu E, et al. Does it matter what you call it? A randomized trial of language used to describe palliative care services. *Support Care Cancer* 2013;21:3411-9.
8. Fadul N, Elsayem A, Palmer JL, et al. Supportive versus palliative care: What's in a name?: A survey of medical oncologists and midlevel providers at a comprehensive cancer center. *Cancer* 2009;115:2013-21.
9. Dalal S, Palla S, Hui D, et al. Association between a name change from palliative to supportive care and the timing of patient referrals at a comprehensive cancer center. *Oncologist* 2011;16:105-11.
10. Parikh RB, Kirch RA, Smith TJ, et al. Early specialty palliative care — translating data in oncology into practice. *N Engl J Med* 2013;369:2347-51.

Affiliation: Department of Family Medicine, Carolinas HealthCare System, Charlotte, NC


A credit to you!

CMA members can earn CME credits for reading CMAJ articles

The *CMAJ* is committed to helping meet your accredited learning goals. CMA members can earn CME credits simply by reading the *CMAJ*. Selected *CMAJ* articles are eligible for Mainpro-M1 credits from The College of Family Physicians of Canada (CFPC). Members can also obtain Mainpro-M2 credits or MOC (Section 2) credits from Royal College of Physicians and Surgeons of Canada (RCPSC) for reading any article in *CMAJ*.

Learn more

www.cmaj.ca/site/cme



Dr. Stephen Pomedli
Family physician
Toronto, Ont.

CMAJ
2986