

Appendix 2 (as supplied by the authors): Price increase per pack of cigarettes from baseline as a predictor of cessation outcomes at 6 months and 1 year

Outcome	Risk associated with a \$1 increase per 20 cigarettes, RR (95% CI)*	
	Unadjusted	Adjusted†
Attempt to quit		
6 mo	1.08 (1.02–1.14)	1.06 (0.98–1.15)
1 yr	1.09 (1.04–1.15)	1.04 (0.98–1.11)
30-day cessation		
6 mo	1.08 (0.99–1.17)	1.06 (0.98–1.15)
1 yr	1.10 (0.99–1.22)	1.07 (0.98–1.15)
Long-term cessation		
1 yr	1.11 (0.97–1.27)	1.12 (1.01–1.25)
Note: CI = confidence interval, RR = relative risk. *Sample size for unadjusted analyses was 881; sample size for adjusted analyses was 721. †Adjusted for all covariates listed in Table 1.		

Appendix to: Mecredy GC, Diemert LM, Callaghan RC, et al. Association between use of contraband tobacco and smoking cessation outcomes: a population-based cohort study. *CMAJ* 2013; DOI:10.1503/cmaj.111861.

Copyright © 2013 The Author(s) or their employer(s).

To receive this resource in an accessible format, please contact us at cmajgroup@cmaj.ca.