

Appendix 1: References

1. Schulman S, Lindmarker P, Holmstrom M, et al. Post-thrombotic syndrome, recurrence, and death 10 years after the first episode of venous thromboembolism treated with warfarin for 6 weeks or 6 months. *J Thromb Haemost* 2006;4:734-42.
2. de Wolf MA, Wittens CH, Kahn SR. *Phlebology* 2012;27(Suppl 1):85-94.
3. Kahn SR, Hirsch A, Shrier I. Effect of postthrombotic syndrome on health-related quality of life after deep venous thrombosis. *Arch Intern Med* 2002;162:1144-8.
4. Kolbach DN, Sandbrink MW, Hamulyak K, et al. Non-pharmaceutical measures for prevention of post-thrombotic syndrome. *Cochrane Database Syst Rev* 2004;(1)CD004174.
5. Kahn SR, Shbakio H, Shapiro S, et al. Effectiveness of compression stockings to prevent the post-thrombotic syndrome (the SOX Trial and Biol-SOX biomarker substudy): a randomized controlled trial. *BMC Cardiovasc Disord* 2007;7:21.
6. Vazquez SR, Kahn SR. Advances in the diagnosis and management of postthrombotic syndrome. *Best Pract Res Clin Haematol* 2012;25:391-402.