

Supplementary Table 3: Logistic regression for the odds of cancers diagnosed at higher stage during SOE vs. before SOE, adjusting for time trend and seasonality

Cancer	Ordinal logistic regression		Stage III, IV vs Stage I, II		Stage IV vs Stage I, II, III	
	OR (95% CI)	p-val	OR (95% CI)	p-val	OR (95% CI)	p-val
Bladder	1.04 (0.77, 1.41)	0.80	0.71 (0.43, 1.16)	0.17	0.61 (0.30, 1.22)	0.16
Breast	1.15 (0.96, 1.38)	0.12	1.14 (0.89, 1.46)	0.29	1.34 (0.93, 1.92)	0.11
Cervix	0.89 (0.45, 1.78)	0.75	0.73 (0.27, 2.02)	0.54	2.11 (0.45, 9.85)	0.34
Colorectal	1.17 (0.96, 1.43)	0.11	1.13 (0.90, 1.41)	0.30	1.12 (0.87, 1.44)	0.36
Kidney	1.11 (0.77, 1.59)	0.59	0.98 (0.66, 1.44)	0.91	1.24 (0.73, 2.11)	0.43
Lung	0.97 (0.81, 1.16)	0.71	0.98 (0.79, 1.20)	0.82	0.98 (0.81, 1.18)	0.82
Melanoma	1.35 (0.93, 1.95)	0.11	1.42 (0.90, 2.26)	0.14	1.25 (0.57, 2.74)	0.58
NHL	1.11 (0.82, 1.51)	0.49	0.95 (0.67, 1.35)	0.77	1.23 (0.88, 1.72)	0.22
Prostate	1.49 (1.22, 1.81)	<.001	1.58 (1.27, 1.96)	<.001	1.42 (1.09, 1.86)	0.01
Uterus	1.14 (0.76, 1.71)	0.51	1.16 (0.74, 1.81)	0.52	1.06 (0.56, 2.00)	0.87