

Appendix 2 (as supplied by authors): Respiratory infections with fever greater than 38° among mothers and infants,* by influenza period and vaccine group						
Group; outcome	No influenza virus circulating, Sept. 2004–Jan. 2005			Influenza virus circulating, Feb. 2005–Oct. 2005		
	Control	Influenza	<i>p</i> value	Control	Influenza	<i>p</i> value
Mothers, no.	107	104		59	64	
Episodes of respiratory illness with fever, no.	17	14		16	5	
Person-months	654	649		422	440	
Incidence/100 person-months (95% CI)	2.60 (1.51 to 4.16)	2.16 (1.18 to 3.62)		3.79 (2.17 to 6.16)	1.14 (0.37 to 2.65)	
Incidence difference (95% CI)	–0.44 (–2.12 to 1.23)			–2.66 (–4.76 to –0.55)		
Incidence rate ratio (95% CI)	0.83 (0.41 to 1.68)			0.6	0.30 (0.11 to 0.82)	0.01
Infants, no.	109	102		57	52	
Episodes of respiratory illness with fever, no.	20	22		57	34	
Person-months	278	265		592	613	
Incidence/100 person-months (95% CI)	7.19 (4.39 to 11.11)	8.30 (5.20 to 12.57)		9.63 (7.29 to 12.47)	5.55 (3.84 to 7.75)	
Incidence difference (95% CI)	1.11(–3.58 to 5.80)			–4.08	(–7.2 to –0.96)	
Incidence rate ratio (95% CI)	1.15 (0.63 to 2.11)			0.7	0.58 (0.38 to 0.88)	0.01
*All mothers and infants with morbidity data included.						