

Appendix 2: Use of contraception among women aged 40–44 years¹⁻³			
Method of contraception	% of women		
	United States 2006–2008	United Kingdom* 2008–2009	Canada*† 2006
Female sterilization	50.2	18	16.4
Male sterilization	19.6	28	19.6
Oral contraceptive	11.1	10	17.1
IUD	4.2	12	4.6
Condom	8.8	21	42.5
DMPA	1.1	2	1
Periodic abstinence	0.6	4	9
Withdrawal	3	6	8.1

Note: DMPA = depot medroxyprogesterone acetate, IUD = intrauterine device.
 *Respondents could choose more than 1 answer.
 †Canadian data were for women over 40; no upper age limit was specified.

References

1. Mosher WD, Jones J. Use of contraception in the United States: 1982–2008. *Vital Health Stat 23* 2010;Aug:1-44.
2. Lader D. *Contraception and sexual health, 2008–9*. London (UK): Office for National Statistics; 2009. Available: www.ons.gov.uk/ons/rel/lifestyles/contraception-and-sexual-health/2008-09/index.html (accessed 2012 June 26).
3. Black A, Yang Q, Wu Wen S, et al. Contraceptive use among Canadian women of reproductive age: results of a national survey. *J Obstet Gynaecol Can* 2009;31:627-40.