

1 APPENDIX 2 (as submitted by the authors): Supplementary tables

2

3 Supplementary Table 1. Life table for the cohort.

Years	Number at risk	TKR	Censored	Annual Incidence (%)	Cumulative Incidence (%)	Cumulative Incidence 95% CI	
						Lower	Upper
0 to 1	643	0	15	0.0	0.0	-	-
1 to 2	628	2	47	0.3	0.3	0.0	1.3
2 to 3	579	4	54	0.8	1.1	0.5	2.3
3 to 4	521	9	40	1.7	2.8	1.7	4.7
4 to 5	472	10	39	2.2	5.0	3.4	7.3
5 to 6	423	5	59	1.2	6.2	4.4	8.8
6 to 7	359	10	44	2.8	9.0	6.6	12.1
7 to 8	305	12	48	3.9	12.9	9.9	16.7
8 to 9	245	14	27	5.2	18.1	14.4	22.7
9 to 10	204	6	31	2.6	20.7	16.6	25.7
10 to 11	167	5	43	2.8	23.5	19.0	28.9
11 to 12	119	8	20	5.6	29.1	23.6	35.5
12 to 13	91	7	25	6.3	35.4	28.9	42.9
13 to 14	59	7	15	8.8	44.2	36.1	53.2
14 to 15	37	2	22	4.3	48.5	39.3	58.6
15 to 16	13	1	7	5.4	53.9	41.3	67.5
16 to 17	5	1	4	15.4	69.3	43.7	91.2

4 *Abbreviations:* CI = confidence intervals; TKR = total knee replacement

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24 **Supplementary Table 2. Life table divided by radiographic disease severity (i.e. Kellgren &**
 25 **Lawrence grade).**
 26

Years	Number at risk	TKR	Censored	Annual Incidence (%)	Cumulative Incidence (%)	Cumulative Incidence 95% CI	
						Lower	Upper
Mild-to-moderate (i.e. KL 2 or below)							
0 to 1	260	0	8	0.0	0.0	-	-
1 to 2	252	0	19	0.0	0.0	-	-
2 to 3	233	1	29	0.5	0.5	0.1	3.2
3 to 4	203	0	20	0.0	0.5	0.1	3.2
4 to 5	183	2	19	1.1	1.6	0.5	5.0
5 to 6	162	0	33	0.0	1.6	0.5	5.0
6 to 7	129	1	19	0.8	2.4	0.9	6.5
7 to 8	109	5	24	5.1	7.5	3.9	14.2
8 to 9	80	3	11	3.7	11.2	6.4	19.3
9 to 10	66	1	14	1.5	12.7	7.4	21.4
10 to 11	51	1	19	2.1	14.8	8.7	24.7
11 to 12	31	2	6	6.1	20.9	12.2	34.4
12 to 13	23	0	8	0.0	20.9	12.2	34.4
13 to 14	15	0	4	0.0	20.9	12.2	34.4
14 to 15	11	0	9	0.0	20.9	12.2	34.4
15 to 16	2	0	1	0.0	20.9	12.2	34.4
16 to 17	1	0	1	0.0	20.9	12.2	34.4
Moderate-to-severe (i.e. KL grade 3 or 4)							
0 to 1	383	0	7	0.0	0.0	-	-
1 to 2	376	2	28	0.6	0.6	0.1	2.2
2 to 3	346	3	25	0.9	1.5	0.6	3.4
3 to 4	318	9	20	2.8	4.3	2.6	7.2
4 to 5	289	8	20	2.8	7.1	4.7	10.6
5 to 6	261	5	26	1.8	8.9	6.2	12.8
6 to 7	230	9	25	3.8	12.7	9.3	17.3
7 to 8	196	7	24	3.3	16.0	12.1	21.1
8 to 9	165	11	16	5.9	21.9	17.1	27.8
9 to 10	138	5	17	3.0	24.9	19.7	31.3
10 to 11	116	4	24	2.9	27.8	22.2	34.6
11 to 12	88	6	14	5.4	33.2	26.7	40.8
12 to 13	68	7	17	7.8	41.0	33.3	49.8
13 to 14	44	7	11	10.8	51.8	42.3	61.9
14 to 15	26	2	13	4.9	56.7	46.1	67.8
15 to 16	11	1	6	5.4	62.1	48.7	75.7
16 to 17	4	1	3	15.2	77.3	51.2	95.3

27 *Abbreviations:* CI = confidence intervals; TKR = total knee replacement

28

29

30 **Supplementary Table 3. Life table divided by sex.**

Years	Number at risk	TKR	Censored	Annual Incidence (%)	Cumulative Incidence (%)	Cumulative Incidence 95% CI	
						Lower	Upper
Male							
0 to 1	498	0	12	0.0	0.0	-	-
1 to 2	486	0	39	0.0	0.0	-	-
2 to 3	447	4	46	0.9	0.9	0.4	2.5
3 to 4	397	5	32	1.3	2.2	1.2	4.3
4 to 5	360	8	24	2.3	4.5	2.8	7.2
5 to 6	328	3	46	0.9	5.4	3.5	8.3
6 to 7	279	7	38	2.6	8.0	5.5	11.5
7 to 8	234	9	34	3.8	11.8	8.6	16.1
8 to 9	191	9	20	4.4	16.2	12.2	21.3
9 to 10	162	2	25	1.1	17.3	13.1	22.6
10 to 11	135	5	35	3.5	20.8	16.0	26.9
11 to 12	95	6	17	5.5	26.3	20.4	33.6
12 to 13	72	6	20	7.2	33.5	26.1	42.2
13 to 14	46	6	12	9.9	43.4	34.1	54.1
14 to 15	28	1	17	2.9	46.3	36.1	57.9
15 to 16	10	0	6	0.0	46.3	36.1	57.9
16 to 17	4	1	3	21.5	67.8	36.6	94.0
Female							
0 to 1	145	0	3	0.0	0.0	-	-
1 to 2	142	2	8	1.4	1.4	0.4	5.7
2 to 3	132	0	8	0.0	1.4	0.4	5.7
3 to 4	124	4	8	3.3	4.7	2.2	10.3
4 to 5	112	2	15	1.9	6.6	3.3	12.7
5 to 6	95	2	13	2.1	8.7	4.7	15.6
6 to 7	80	3	6	3.5	12.2	7.2	20.3
7 to 8	71	3	14	4.1	16.3	10.2	25.7
8 to 9	54	5	7	8.3	24.6	16.4	36.0
9 to 10	42	4	6	7.8	32.4	22.5	45.1
10 to 11	32	0	8	0.0	32.4	22.5	45.1
11 to 12	24	2	3	5.9	38.3	26.9	52.7
12 to 13	19	1	5	3.8	42.1	29.5	57.4
13 to 14	13	1	3	5.0	47.1	32.8	64.1
14 to 15	9	1	5	8.2	55.3	37.0	75.4
15 to 16	3	1	1	17.9	73.2	43.3	95.3
16 to 17	1	0	1	0.0	73.2	43.3	95.3

Abbreviations: CI = confidence intervals; TKR = total knee replacement

31
32
33
34
35

36 **Supplementary Table 4. Sensitivity Analysis. Cox regression hazard ratio estimates for**
 37 **independent predictors of conversion from HTO to TKR when including only a single limb**
 38 **from each patient (n=556).**

Predictor	Hazard Ratio (95% CIs)
Radiographic severity (Kellgren & Lawrence [KL] grade)	
Mild-to-moderate*	Reference
Moderate-to-severe*	2.47 (1.26 to 4.84)
Pre-operative alignment (Mechanical axis angle [degrees])	1.05 (0.97 to 1.14)
Correction Size (mm)	1.03 (0.93 to 1.14)
KOOS Pain Score (per 10 points) ^a	0.77 (0.68 to 0.89)
Sex	
Male	Reference
Female	1.60 (0.98 to 2.61)
Age (per 10 years)	1.40 (1.04 to 1.88)
Body mass index (per 5 kg/m ²)	1.26 (1.04 to 1.52)
Year of surgery	1.01 (0.92 to 1.11)

39 * Mild-to-moderate = KL grade 2 or less, moderate-to-severe = KL grade 3 or 4

40 ^a KOOS = Knee injury and Osteoarthritis Outcome Score – 0 indicates extreme knee symptoms; 100 indicates no
 41 knee symptoms.

42 *Abbreviations:* CI = confidence intervals

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71

72 **Supplementary Table 5. Post hoc Analysis. Cox regression hazard ratio estimates for**
 73 **independent predictors of conversion from HTO to TKR (n=643) stratified by surgeon.**

Predictor	Hazard Ratio (95% CIs)
Radiographic severity (Kellgren & Lawrence [KL] grade)	
Mild-to-moderate*	Reference
Moderate-to-severe*	2.07 (1.19 to 3.62)
Pre-operative alignment (Mechanical axis angle [degrees])	1.05 (0.98 to 1.11)
Correction Size (mm)	1.04 (0.96 to 1.13)
KOOS Pain Score (per 10 points) ^a	0.86 (0.75 to 0.97)
Sex	
Male	Reference
Female	1.66 (1.06 to 2.60)
Age (per 10 years)	1.44 (1.12 to 1.85)
Body mass index (per 5 kg/m ²)	1.34 (1.14 to 1.58)
Year of surgery	1.07 (0.97 to 1.18)

74 The model was stratified by surgeon and the variance was adjusted for staged bilateral osteotomies using robust
 75 sandwich estimators.

76 * Mild-to-moderate = KL grade 2 or less, moderate-to-severe = KL grade 3 or 4

77 ^aKOOS = Knee injury and Osteoarthritis Outcome Score – 0 indicates extreme knee symptoms; 100 indicates no
 78 knee symptoms.

79 *Abbreviations:* CI = confidence intervals

80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103

104 **Supplementary Table 6. Post hoc Analysis. Cumulative incidence of conversion from HTO**
 105 **to TKR in patients categorized by size of surgical correction angle.**
 106

Postoperative Alignment* (Mechanical Axis Angle)	Number of knees	5-year Cumulative incidence	10-year Cumulative incidence
0° to 3° valgus (i.e. neutral)	319	1.9 (0.8 to 4.6)	13.9 (9.3 to 20.5)
< 0° varus	205	9.2 (5.6 to 15.1)	30.0 (22.1 to 40.0)
> 3° valgus	119	6.3 (2.9 to 13.6)	23.0 (14.0 to 36.4)

107 * Values are presented as percentages (with 95% confidence intervals)

108 Correction angle was assessed using the first postoperative full-limb standing radiographs and the mechanical axis
 109 angle. We defined correction angle by 3 categories based on previous literature^{1,2}.

110 *Abbreviations:* CI = confidence intervals

111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142

143 **Supplementary Table 7. Post hoc Analysis. Cox regression hazard ratio estimates for**
 144 **independent predictors of conversion from HTO to TKR (n=643) that includes correction**
 145 **angle.**

Predictor	Hazard Ratio (95% CIs)
Radiographic severity (Kellgren & Lawrence [KL] grade)	
Mild-to-moderate*	Reference
Moderate-to-severe*	2.07 (1.17 to 3.64)
Pre-operative alignment (Mechanical axis angle [degrees])	1.05 (0.99 to 1.11)
Correction angle (degrees) ^a	
0° to 3° valgus (i.e. neutral)	Reference
< 0° varus	1.87 (1.18 to 3.00)
> 3° valgus	1.40 (0.80 to 2.46)
KOOS Pain Score (per 10 points) ^b	0.87 (0.77 to 0.98)
Sex	
Male	Reference
Female	1.68 (1.07 to 2.62)
Age (per 10 years)	1.59 (1.25 to 2.03)
Body mass index (per 5 kg/m ²)	1.30 (1.11 to 1.53)
Year of surgery	1.04 (0.95 to 1.13)

146 The variance was adjusted for staged bilateral osteotomies using robust sandwich estimators.

147 The predictor for correction size was substituted for correction angle as the variables are multicollinear.

148 * Mild-to-moderate = KL grade 2 or less, moderate-to-severe = KL grade 3 or 4

149 ^a Correction angle was assessed using the first postoperative full-limb standing radiographs and the mechanical axis
 150 angle. We defined correction angle by 3 categories based on previous literature^{1,2}.

151 ^b KOOS = Knee injury and Osteoarthritis Outcome Score – 0 indicates extreme knee symptoms; 100 indicates no
 152 knee symptoms.

153 *Abbreviations:* CI = confidence intervals

154

155

156

157

158

159

160 References

161

162 1. Fowler PJ, Tan JL, Brown GA. Medial Opening Wedge High Tibial Osteotomy: How I
 163 Do It. *Operative Techniques in Sports Medicine*. 2012;20(1):87-92.

164 2. Hui C, Thompson SR, Giffin JR. Chapter 103 - Knee Arthritis. In: Miller MD, Thompson
 165 SR, editors. *DeLee & Drez's Orthopaedic Sports Medicine Textbook*,. 4th ed. New York, NY:
 166 Elsevier; 2014. p. 1227-42.

167

168